

EOS, Transactions, American Geophysical Union

1.0

Vol. 64, No. 12, Pages 113 - 120

March 22, M

Particles and Fields— Magnetosphere

5120 Interactions between solar wind and magnetosphere
COMPARISON OF 33-3 POLAR CAP POTENTIAL DROPS WITH THE
INTERPLANETARY MAGNETIC FIELD AND MODELS OF MAGNETOPAUSE RECONNECTION

J. R. Wysam (Space Scences Laboratory, University of California, Berkeley, California, 94730, R. B. Tower, and F. S. Mozer
Measurements of the cross polar cap electric potential, by the double
probe electric field experiment aboard 33-3, from fifty-five critis in the
data-in-dusk plane are compared to the reconnection electric fields
predicted by a variety of models, both the ordered and experimental. The
purpose of these comparations is to understand the extent to which nonreconnection contributions to the polar cap potential must be included, to
delarating the time exposus of the polar cap potential in the terminal of the reconnection ruses, and to determine the efficiency and inturvation levels
of the reconnection process.

reconnection ruses, and to determine the efficiency and subration levels of the reconnection process.

It is found that III After several boars of northward interplanetary magnetic field, the cross polar cap potential declinas to progressively lower values than those after one hour of northward laterplanetary magnetic field. This suggests that it requises several hours for the tomospheric polar cap potential to respond to the 'terming off' or 'turning doesn' of the reconnection process [27] The decay of the polar cap potential is used to demonstrate that constitutions to the polar cap potential in associated with the reconnection process cut be limbed to less than 20 killowits. It is shown that contributions to the polar cap potential that scale with the dynamic pressure of the after which are limited to less than one killowit. It is shown that contributions from interplanetary magnetic field parameters over the four hours previous to the measurement. The weighing functions have the four hours previous to the measurement. The weighing functions have the four hours previous to the measurement. The weighing functions have the four hours previous to the measurement of the down dask component of the interplanetary electic field less than eboot 0.5 meViranter, the measurem polar cap potantial is contistent with reconnection of all the laterphenetary magnetic field from their passessary magnetic field for incident on a 100% with frontistic magnetics. For ent on a 10/Rs wide frontsis

magnetic flux incident on a 10-ftg wide frontiside magneticpasses. For larger values of the dewn dark component of the interplaneary electric field, the proportion of field lines reconnecting is less, indicating servertion at the reconnection process.

The above, results are obtained when the extended response time of the magnetosphere to changes in interplaneary parameters is considered. They are independent of the desired reconnection model essented, and they are quite different from the viscous contribution, saturation lively of reconnection, and reconnection efficiencies inferent from comparisons of potar cap potentials to digital hour averages of interplaneary parameters.

J. Geophys. Res., Blue, Peper 3AQ277-

J. Geophys. Res., Blue, Paper 1AC177.

5775 (interactions Between Nagnetosphere and Counc. Roys.)

SITE (interactions Decrease in Satural's Rights) COUNCE ROYs.)

SITE (interactions Decrease in Satural's Rights) COUNCE ROYs.)

RICLER CASCAGES IN SATURAL'S RIGHTS COUNCE ROYs.)

RICHE HARCH MANGETCHERGE

John F. Cooper (Enrice Fermi Institute, University of Chicago, 633 st. 56th Street, Chicago, 111 inch 80637)

The nearly equalorial trajectory of the Ploteer II spacecraft through Saturn's high energy proton radiation belts and under the smith Association for proceeding the Ploteer II spacecraft through Saturn's high energy proton radiation belts and under the smithest and the Police II in the 230 legan and supportantity to study the radial dependence of the satural proton interactions in the rings. Sensitive proton injection in the reduction belts by neutron bate-decay, assentespheric diffusion, and absorption by planetary rings and stabilities. Maximum trapped proton interaction in the reduction belts are compared with calculated inheaptites and found consistent with trapped grade in the processing of the compared with passing planets of the form of the spectra intended from integral mental energy spectra intended fr

10H-BEAM-DRIVEN ELECTROSTATIC ION CYCLOTRON
INSTABLLITIES
A. Hiura (Geophysics Research Laboratory, University
of Tokyo, Bunkyo-ku, Tokyo, 113 Japan), H. Chuda and
M. Ashour-bhdaila
We present results of numerical simulations of
slockrostatic ion cyclotron instabilities driven by
an ion beam parallel to the augmoric field. For
beam speed exceeding the thermal speed of the background ions and beam temperature such lover than the
background ion temperature, it is found that the
instability results in strong perpendicular hosting
and slowing down of the parallel drift of beam ions,
leading to the saturation of the instability.
Applications to plasma heating and space plasma
physics are discussed. (ion beams, ion electroquality waves, arroral instabilities).

5753 Plasma instabilities.

5755 Plagma instabilities DRIFT MAYE MODEL FOR GEOMAGNETIC PULSATIONS IN A HIGH #

FLARMA
V.L. Fatai, P.H. Mg (Capatiment of Physics, University
of Danovar, Danovar, CO 80208) and L.J. Cachill, Jr.
(Space Science Center, University of Minnesota,
Minneapolia, NM 55455)
A dispersion relation for the stability of the
coupled drift compressional drift mirror and shoar
Alfvéa wavas in the magnotospheric plasses is enslysed
by numerical method. The smalysis is suitable for the
storm-time plassa conditions in the magnatosphero which
usually has gradients in plassa demaity, temperature,
of high \$ which is characteristic of storm-time plasse
is included in the model. For a given appropriate aat
of plassa paraseters, and wave parameters of a mode,
relative wave amplitudes are calculated from the model.
A comparison is made of model generated and observed
relative wave amplitudes by wing two sample wave relative wave amplitudes by using two sample wave events observed by Explorer 45 during August 4-5, 1972 supports etcm. The quantizative amelysis shows good agreement between theory and observations for a gradient-driven alfred-like instability showever growth rates are very small for this mode. The anisotropy-driven drift alreor instability has large growth rates are very small for this mode. The anisotropy-driven drift alreor instability has large growth rates, but does not show good correlation between theory and observations of raintive wave amplitudes. (Drift wave instabilities, geomagnetic polastions).

J. Goophys. Res., Shue, Paper JAD400

Planetology

0510 Almonyhurus of Planets
ELECTRINI PRECIPITATION AND RELATED ARROWSHY OF THE
JOHN THERMSCHINERS AND TOROUGHERS
J. HUNDER Walte, Jr. Illjace Soience Laboratory,
NAUA/Moreheal Hyster Flight Center, AL 385121, 7. 1.
Cravens, Jr. Koryra, A. P. Nogy, S. K. Akrays, and L.
Cravens, Jr. Koryra, A. P. Nogy, S. K. Akrays, and L.
Chan

MANA/Harshail figure Figure Center, as Arrays, and L. Cravens, ... Knayra, A. P. Nayv, S. E. Arrays, and L. Cravens, ... Knayra, A. P. Nayv, S. E. Arrays, and L. Cravens, ... Knayra, A. P. Nayv, S. E. Arrays, and L. Cravens, ... Characteristics and therman hand remainded at high latitudes indicated the many form a. C. the detail no morely fluor of about Merical and the many form and the many figure of the control of the co and vibrational heating of M. from predictable processes appear to play a distral role is described the structure of the high-latitude longshers. Theoretical fits to the Voyager radio consisters electron density profiles at high latitudes of the local structure of the latitude structure of the local structure which reaches 3000 K in the local structure of the local

Londaphara. (Auplter, aurora, aboals ny Londaphara.)

J. Geophys. Ras., Blua, Paper 3A0222

6599 General or Miscellaneous
ENERGETIC ION LOSSES MEAR IO'S GREIT
A. F. Chang (Dept. of Physics and Astronomy, August Miniversity, Piscetaway, H.J. 0886) C. S. Meligned University, Piscetaway, H.J. 0886) C. S. Meligned University, Piscetaway, H.J. 08869 C. S. Meligned Trom the Low Energy Charged Particle (LEC) engineers of the Wood of the Company of the Company of the Meligned Company of the Wood of the Company of the Company of the Wood of the Company of the Wood of the Company of th

3-D Model of the Lithosphere-Asthenosphere Boundary for a Propagating Rift

Phillip D. Milholland and Frederick K. Duennebier

Hawaii Institute of Geophysics University of Hawaii at Manoa Ionolulu, Hawaii

A first-order approximation of the litho-sphere-asthenosphere boundary of a propa-gating rift model is shown in Figures 1 and 2. (Figure 1 is a reduced version of the cover figure. All bracketed letters in the text correspond to letters in Figure 2.) The propagating rift model explains tectonic patterns of some oceanic regions that have puzzled earth scientists for some time. According to this model, oceanic spreading centers can propa-gate through older crust, transferring portions of crust from one plate to another.

The two images in Figure 1 form a 3-D ste-

reo pair of the lithosphere-asthenosphere boundary at a propagating rift. To view the 5-D model use a stereoscope on Figure 1; alternatively, stare at Figure 1 or the cover figure and let your eyes soft-focus until a 3-D image appears between the two printed images (believe us, it works). The model is based on the Galapagos propagator at 95.5°W, viewed from above and toward the southeast. The 'V'-shaped wedge of litho-sphere between the pseudofaults is formed by extension of the eastern rift toward the viewer. As this rift propagates, spreading stops on the dying western rift, leaving a failed rift. Only two plates are present in the model, and depth to the asthenosphere depends only on

The stippled region (A) represents a cross section of the lithosphere and shows how the lithosphere thickens away from the rift. Depth to the ocean bottom (top boundary of A) is calculated by using the equations and assumptions of Schater and Francheteau [1970] for a spreading center that spreads at a rate of 26 mm/yr. Depth is 2.1 km at the spreading center and increases to 3.1 km on the far ht. Lithospheric thickness is calculated by using the equation of Yoshi et al. [1976]

 $L = 7.49(T)^{0.5}$

where L is the thickness of the lithosphere in kilometers, and T is the age in millions of years. Depth to the lithosphere-asthenosphere boundary (bottom edge of A) is the sum of the ocean depth and lithospheric thickness. Thermal and isostatic adjustments that occur at the tip of the propagator and other boundaries have not been incorporated into the model and would tend to smooth the discontinuities and sharp boundaries. Depth is assumed to be a function of age only.

General configuration of the model corresponds to the 95.5°W Galapagos propagator [Her et al., 1980], except that the initial rifts of this model are offset by a 20-km-long transform fault (B2). Fracture zones (B1, B2-B3) associated with this transform fault terminate at the ends of the pseudofaults (C.1, C.2). The southern fracture zone (B2-B3) is composed of two sections: the fracture zone that existed when the transform was active (B3) and the original transform fault that was locked in place when rift propagation com-menced (B2). As Hoy [1977] pointed out, and as can be seen clearly here, the sense of vertical offset changes along the southern fracture zone instead of monotonically decreasing, as it does along a normal fracture zone. Intersections of the pseudofaults with the fracture zones mark the locations of lithosphere formed at the initiation of propagation. Propagation began at the initial transform fault and has extended the length of the eastern rift (D1) at a constant rate along a different azimuth. Extension of the propagating rift or propagator (D2) proceeds at the expense of the dying rift (D3). As spreading is trans-

Ahoy! Sail Back into Baltimore

SAVE MONEY... Register perfore May 11
Tickets still available for Section luncheon and pesident's bin Fly United call toll free 800-521 fogsion Michigan residents 100/482-07 or Deltais Call toll page 800-521 fogsion Michigan residents 100/482-07 call toll page 800-521 fogsion Michigan residents 100/482-07 or Deltais 100 fogsion Michigan residents 100/482-07 call toll page 800-521 fogsion Michigan

Reserve early for preferred meeting hotel

Housing Deadline: April 25

ferred to the propagator, a portion of the dy-ing rift is frozen into the southern plate and becomes a failed rift (D4). Orientation of the failed rift is controlled by the spreading rate and the extensional growth rate of the propa-gator. There are only two rigid plates in this model at all times.

Pseudofaults (C1, C2) stand out as large depth discontinuities in the lithosphere as-thenosphere boundary: these discontinuities reflect differences in age between new lithosphere formed at the propagator tip and old lithosphere through which the propagator extends. Vertical offset along the pseudofaults decreases away from the propagator tip because the difference in thickness decreases as both sections of the lithosphere age and because the propagator is extending through progressively older and thicker lithosphere. Note that no horizontal displacement occurs along the pseudofault boundaries, although vertical motion should result from differential aging across the boundaries.

The transform fault (E), which connects the propagator tip to the intersection of the dying and failed rifts, differs in important ways from the classical transform fault of a normal rift system. Material on both sides of the transform fault is formed by the same rift; consequently, there is no sharp age or thermal difference across the boundary. Also, the location of the transform fault is not fixed but moves in the same direction and at the same rate as rift propagation. Its length changes if the propagator is growing at a dif-ferent azimuth from the dying rift. Note that what would be fracture zones (inactive portions of the transform) in a normal transform fault system are manifested by the northern pseudofault (C1) and by the zone between the southern pseudofault (C2) and the failed rift (D4). All interplate horizontal displacement occurs in the region between the propagator tip and the dying rift and is recognized by the change in azimuth between the dying and failed rifts. Hey et al. [1980] suggest that the transform fault of a propagating rift may be a zone of shear deformation rather than a localized fault boundary.

The existence of propagators in several areas has been postulated [Hey, 1977; Hey et al., 1980; Hey and Wilson, 1980; Delaney et al. 1981). Propagators, and the causes and consequences of propagation, will be one of the most interesting topics in plate tectonics to be studied during this decade.

Acknowledgments

This work is funded under National Science Foundation grant OCE-78-19816. The plotting software was written by J. Tuthill and D. Cuddy. The authors thank Jim Kel-logg, Rita Pujalet, and Brian Taylor for critically reviewing this paper. This is Hawaii Institute of Geophysics Contribution 1358.

Delaney, J. R., H. P. Johnson, and J. L. Kar-sten, The Juan de Fuca Ridge—Hot spot— Propagating rift system: New tectonic, geochemical and magnetic data, J. Geophys. Res., 86 (B12), 11,747-11,750, 1981.

Hey, R. N., A new class of 'pseudofaults' and their bearing on plate tectonics: A propagating rift model, Earth Planet. Sci. Lett., 37, 321-325, 1977. Hey, R. N., and D. S. Wilson, Propagating rifus—The motion picture, Eos Trans. AGU,

61, 1104–1105, 1980. Hey, R. N., F. K. Duennebier, and W. J. Morgan, Propagating rifts on midocean ridges, J. Geophys. Res., 85, 3647-3658, 1980.
Sciater, J. G., and J. Franchetau, The impli-

cations of terrestrial heat flow observations on current tectonic and geochemical models of the crust and upper mantle of the

> 1983 **AGU**

Spring

Fig. 1. A 3-D stereo pair of the lithosphere-asthenosphere boundary at a propagating rift. See text for details.




Fig. 2. (A) Lithosphere—The upper boundary is the depth below sea level (2.1-3.1 km), and the lower boundary is the lithosphere-asthenosphere boundary (2.1-21 km). (B) Fracture Zones—B1 and B3 are fracture zones that form from an original transform fault, B2, that is locked in place when rift propagation commences. (C) Pseudofaults—Pseudofaults mark the boundary of lithosphere formed by the propagator. The large vertical offset is a consequence of the large age difference across the boundary. (D) Rifts—D1 is an original rift from which the propagating rift or propagator, D2, arises. D3 is the dying rift which becomes inactive and forms the failed rift, D4, as the propagator usurps the spreading. (E) Transform Fault—The transform fault lies between the propagator tip and the end of the dying rift. It is significantly different from a classical transform fault in that material on both sides of the fault is formed by the same rift, there is no significant age difference across the fault, no fracture zones develop, and the fault must be repositioned at the same across the fault, no fracture zones develop, and the fault must be repositioned at the same rate as the extensional growth rate of the propagator. Scale in kilometers.

earth, R. Astron. Soc., Geophys. J., 20, 509-Yoshi, T., Y. Kono, and K. Ito, Thickening of the oceanic lithosphere, in *The Geophysics of the Pacific Ocean Basin and Its Margin, Geophys. Monogr. Ser.*, vol. 19, edited by G. H. Sutton, M. H. Manghnani, and R. Moberly, pp. 423–430, AGU, Washington, D.C., 1976.

Phillip D. Milholland received his B.S. degree from the University of Washington in 1975 and his M.S. from the University of Hawaii in 1978. His Ph.D. work is on the seismicity of the 95.5°W Galapagos propagating

Frederick K. Duennebler received his B.S. from Trinity College, Hartford, Connecticut, in 1965; he received his M.S. (1968) and Ph.D. (1972) from the University of Hawali. He worked on the Apollo Lunar Seismic Experiment and Viking Martlan Seis-mic Experiments until 1975, when a lack of data drove him to marine

eismology and tectonics,



Sets the groundwork for future study The earth's crustal evolution; the structure, density and homogeneity of the earth's core, its dynamics and thermal evolution are reported on in this essential volume. Current research of geochemical and geophysical evolution dealing with Internal convective motions, plate tectonic patterns, viscosity in the mantle, and the history of the earth is contained in this important edition.

300 pages • Illustrated • \$25.00 30% Member discount

Orders under \$50-must be prepaid American Geophysical Union 2000 Florida Ava., N.W. Washington, D.C. 20009 Call 809-424-2488 toll free 462-6903 (local)

121

Robert M. White, president of the University Corporation for Atmospheric Research (UCAR) and an AGU Fellow, has been elected president of the National Academy of Engineering (NAE). White's 4-year term begins July 1. He succeeds Courtland D. Perkins, who has been NAE president since 1975.

As NAE president, White will serve as vice chairman of the National Research Council (NRC). Frank Press, president of the National Academy of Sciences and former AGU president, is NRC chairman.

A search committee has been established at UCAR to find White's successor. For additional information, write to Thomas Dona-luc, Chairman, Search Committee, Universi-ty Corporation for Atmospheric Research, P.O. Box 3000, Boulder, CO 80307. May 6 is the deadline for applications. UCAR, a con-sortium of 50 universities with doctoral programs in the atmospheric sciences or closely related fields, manages the National Center for Atmospheric Research under contract with the National Science Foundation, UCAR also carries on other activities to promote atmospheric science in the public interest.

New Weather Index

Scientists at the National Oceanic and Atmospheric Administration (NOAA) and the University of Delaware have refined the wind-chill factor, a common measurement of weather discomfort, into a new misery regis-ter called the weather stress index. In addition to the mix of temperature and wind speed data used to calculate wind chill, the recipe for the index adds two new ingredients-humidity and a clash of benchmark statistics—to estimate human reaction to weather conditions. NOAA says that the weather stress index estimates human reaction to weather conditions and that the reaction depends on variations from the 'normal' conditions in the locality involved.

Discomfort criteria for New Orleans, La., and Bismarck, N.D., for example, differ drastically. According to NOAA, when it's the middle of winter and it's -10°C with a relative humidity of 80% and 24 km/h winds, persons in New Orleans would be highly stressed while those in Bismarck wouldn't bat an eye.

NOAA plans to generate daily, weekly, and monthly weather-stress maps of the United

TV Special on Geophysics

Earthquake prediction, earthquake preparation in California and Japan, the theory of plate tectonics, and the causes and effects of earthquakes and volcanoes will be the subjects of a National Geographic television special scheduled to air on public television on April

Among the locations visited by Born of Fire are Iceland, where magma oozes to the surface on the remote island of Heimaey, illustrating the moving crustal plates; the Republic of Djibouti in east Africa, where some scientists believe a new ocean will form as three crustal plates spread apart; and the island of Santorini in the Aegean Sea, where a series of earthquakes and volcanic eruptions some 3,500 years ago destroyed two-thirds of the island and obliterated the city of Akrotiri.

the island and obliterated the city of Akroun.

The special, featuring geologist Robert Ballard of the Woods Hole Oceanographic Institution, is produced by the National Geographic Society and WQED of Pittsburgh with a continuing grant from Gulf Oil Corporation. Check local television listings for time

Geophysical Events

This is a summary of SEAN Bulletin, 8(2), February 28, 1983, a publication of the Smithsonian Institution. The complete Long Valley, Colima, and Langila reports are included; the earthquake report is an excerpt. The complete builtein is available in the microfiche edition of Facas a priceofishe and

microfiche edition of Ees as a microfiche supplement or as a paper reprint. Subscriptions to SEAN Bulletin are also available. For the microliche, order document E83-003 at \$2.50 from AGU Fulfillment, 2000 Florida Avenue, N.W., Washington, DC 20009. For reprints, order SEAN Bulletin (give volume and issue numbers and issue clate) through AGU Separates: \$5.50 for one copy of each issue number for those who do not have a deposit account; \$2 for those who do not have a deposit account; \$2 for those who do; additional copies of each issue number are \$100. For
a subscription, order SEAN Bulletin from
AGU Fulfillment; The price is \$18.00 for 12
monthly issues mailed to a United States adder must be prepald.

122

Volcanic Events

Kilauea (Hawaii): Renewed fountaining and

hemisphere cloud; tiny aerosols recondense above 30 km; unusual sunrises and sunsets. Colima (México): Lava extrusion ended June 1982 but plume emission continues.

Ol Doinyo Lengai (Tanzania): Tephra emission continues; lava flow. Langila (New Britain): Explosions build to 6-

Manam (Bismarck Sea): Rumblings, night glow, increased vapor emissions. Ruapehu (New Zealand): Possibly pre-erup-

tivity; rain-caused debris flows.

Long Valley Caldera, California, USA (37.68°N, 118.86°W). As of early March, an average of 10-30 events per day of magnitude ≥1 continued to occur in the southern part of the caldera in the epicentral area of the major January earthquake swarm (see SEAN Bulletin 7 (12) and 8 (1)). For several months prior to the January swarm the back-ground level of seismicity in the caldera had averaged 8-10 earthquakes of magnitude ≥1 per day. Few larger events were recorded in February, but 5 shocks with magnitudes >3 occurred February 18-19 and a magnitude 4 earthquake was recorded February 24 in the January epicentral region. Heavy snows have severely limited deformation monitoring, but available data suggest that no major changes

have occurred since January.
Information Contact: David Hill, Mail Stop 77, U.S. Geological Survey, 845 Middlefield Rd., Menlo Park, CA 94025 USA.

Colima Volcano, SW Mexico (19.42°N, 103.72 W). A French team reached the northern rim of the summit cone in early December. Storm damage to trails prevented them from reaching the southern side of the cone, so they were unable to see the southern-flank lava flow produced by the eruption that began in Describer 1991 (see SEAA) But that began in December 1981 (see SEAN Bul-letin 7 (1-3)). Only fumarolic activity was observed in the western part of the crater and on the northern flank. Gas of essentially atmospheric composition was emitted at 500°C from the northeastern part of the cone and

from the front of this flow and it may still have been advancing very slowly. The southern flank lava flow appeared to have advanced very little since last observed by Luhr in March 1982. Residents of the area reported that incandescence had ended in

from P. Lowenstein:

crater 2 in January (see last month's SEAN Bulletin) culminated in a rise of the niagma column, with an eruptive phase maximum February 11–16. The February 3-11 buildup of the eruption consisted of approximately hour-long periods of loud, rumbling noises, with deep explosion sounds at 5–30 s intervals. Several times per day at irregular intervals in-dividual explosions produced black, ash-lation columns that rose as much as 3-4 km before being dissipated by the north-westerly winds. Night glow, observed February 3, became more intense during this period. Low strombolian fountaining was visible February 3-5 and 9.

ty, crater 2 simultaneously displayed con-tinuous strombolian fountaining to 100 m and intermittent powerful vulcanian explosions. Most of the vulcanian explosions were laterally directed, while the continuous moderate vapour emissions and the strombolian fountaining were central and vertical, leading to the conclusion that crater 2 may contain two

more or less independent vents, "Seiamic activity consisted of a sub-con-tinuous background of harmonic tremor and strombolian B-type earthquakes." and strombolian in-type carinquakes.

Each individual vulcanian eruption produced large amplitude, low-period explosion events. The most powerful of these

lava flow production on E Rift.

Mt. St. Helens (Washington): Spine added to

February lobe, then extrusion stops; seismicity suggests renewed extrusion by late

Long Valley (California): Seismicity remains elevated, but no new swarms. El Chichón (México): Little change to N

day strombolian-vulcanian event.

tive changes continue. Sakurajima (Japan): Increased explosive ac-

from a vent that had recently extruded a lava flow. Rockfalls occurred several times per day

James Luhr and others visited Colima in mid-January and again in early February. June 1982. Plume emission continued in early 1983 at about the same intensity as a year earlier, but there were no episodic increases

in intensity of plume emission as there had been in early 1982. Information Contact: Jean Louis Che-minée, Laboratoire de Géologie, Ecole Nor-male Supérieure, 46 Rue d'Ulm, 75230 Paris Cedex 05, France: James Luhr, Department of Geology and Geophysics, University of California, Berkeley, CA 94720 USA.

Langila Volcano, New Britain Island, Papua New Guinea (5.53°S, 148.42°E). This report is

"The increased vulcanian activity of

"During the 6 days of maximum activi-

Antarctic Research Serles 36

ISBN 0-87590-186 Geology of the Central Transantarctic Mountains

(1983) Edited by Mort D. Turner & John E. Splettstoesser PUBLISHED IN MINIBOOK FORMAT. Papers 1, 2, and 3, illustrated, 62 pages, \$12.95

Paper 1 - Basement Geology of the Beardmore Glacier Region, J.D. Gunner

Paper 2 - Triassic Vertebrates in the Transintarctic Mountains, Edwin Fl. Colbert Paper 3 — Forms and Facles of Vertebraria in Relation to Gondwana Coal, James M. Schopf

15 papers and 4 USGS reconnaissance maps are planned for this volume. FREE catalog listing all of the ARS volumes in print. Please request physic cience fitles, biological science titles or both catalogs

Call **toIl free 800-424-2488**

in the Washington, D.C. area 462-690 American Geophysical Union 2000 Florida Avenue, N.W. · Washington, D.C. 20009

explosions occurred February 12-13 and 15 at the rate of 2-5 per hour.

"Activity diminished rapidly on February 16 and stopped completely on February 17, but harmonic tremor continued. Weak glow was visible on February 19, and vulcanian explosions occurred on February 19 and 23.

"During the eruption, crater 3 (a separate composite cone 300 m west of crater 2) released only weak, white vapours. However, the volume of emission in creased to moderate or large during the first 10 days of February, the time of the activity buildup at crater 2." Information Contact: P. Lowenstein, Senior

Government Volcanologist, Rabaul Volcano Observatory, P.O. Box 386, Rabaul, Papua New Guinea.

Meteoritic Events

Meteorite fall: Tennessee, USA, January 28; additional fireball observation. Fireballs: North Adantic; Australia (2); Bay of Bengal; Egypt; England; Germany; Italy; Oregon and Washington, USA.

Earthquakes

Date	Time (UT)	Magnitude	Latitude
Feb. 13 06 Feb. 14 03 Feb. 25 18	0140 0636 0320 1822 1214	6 5.6 m _b 0 6.3 M _s * 2 4.8 m.	39,99°N 13,84°N 54,96°N 42,10°N 85,96°N
	-		

Longitude	Depth of Focus	Region
75.10°E	shallow	Southwestern Chin
144.98°E	169 km	Mariana Islands
159.19°W	shallow	Alaska peninsula
21.51°E	10 km	Yugoslavia
135.85°E	83 km	Japan

*Berkeley measurement 6.5 M,

Information Service, U.S. Geological Survey, Stop 967, Denver Federal Center, Box 25046, Denver, CO 80225 USA.

CALL FOR **DELEGATES**

18th General Assembly of IUGG

U.S. scientists planning to attend the 18th General Assembly of IUGG, Hamburg, West Germany, August 15-27, 1983, should notify A. F. Spilhaus, Jr., Secretary, U.S. National Committee, 2000 Florida Avenue, N.W., Washington, D.C. 20009, and indicate in which IUGG association they propose to participate so that they can be officially designated as delegates from the United States.

fransactions, American Geophysical Unio The Weekly Newspaper of Geophysics

For full arricles and meeting reports send on copy of the double-spaced manuscript to Fars AGU at the address below and three copies to one of the editors, or send all four copies to far For news items, send two copies of the doubt spaced manuscript to Em.

Editor-in-Chief: A. F. Spilhaus, Jr.; Editor: Marcel Ackerman, Mary P. Amlerson, Peter & Bell (News), Kevin C. Burke, Bruce Doc, Robe H. Eather (History), Clyde C. Goad, Arnold L Cordon, Louis J. Lanzeroui; Editorial Staff: Gregg Forte -Alunaging Editor, Barbara T. Rd man -- New Writer, Marlo E. Godinez -- New Isern; Kathleen M. Lafferry—Editorial Assistant Production Staff: Patricia Bangett, James 1869 lethwaite, Dae Sung Kim, Lisa Lichtenstein, Vie

Officers of he Union

James A. Van Allen President; Charles L. Drake, President-Elect Leslie H. Meredith, General Secretary: Carl Kishinger, Foreign Stretary: A. F. Spillhaus, Jr., Executive Director Waldo E. Smith, Executive Director Emerkus

For advertising information, contact Robin E. Little, advertising condinator, 202-402-6903.

Copyright 1983 by the American Geophysical Union, Material in this issue may be photocopied by individual scientists for research or date room use. Permission is also granted to use short quotes and figures and tables for publica-tion in scientific books and journals. For pendi-sion for any other uses, contact the AGU Pabli-cations Office.

Views expressed in this publication are those or the authors only and do not necessarily reflect official positions of the American Geophysical Union unicas expressly stated.

Subscription price to members is included in arnual dues (\$20.00 per year). Information on in-attutional subscriptions is available on request Second-class postage paid at Washington, D. G. and at additional mailing offices. Ess, Transctions, American Geophysical Union (ISSN 0096-3941) is published weekly by

American Geophysical Union 2000 Florida Avenue, N.W. Washington, D. C. 20009

Cover. The two images form a three-d mensional stereo pair of the lithosphere asthenosphere boundary for a propagal ing rift. Viewing this bouttdary provided insight into the 5-D structure of the rift insight into the 5-D structure of the rift. system. The propagating rift theory offer explanations for a wide range of geologic cal questions, such as the reorients oceanic rifts or spreading centers, origin of transform faults; origin of continental rifting, generation of high magnetic anomaly zones, and formation of complete magnetic anomaly patterns in areas such as the Juan de Fuca plate. Additional discussion and a description of how to view the model in 8-D is included in the accompanying article. (Figure courtesy of Philipp. D. Milholland and Frederick K. Duenne-

Advanced Techniques for Clay Mineral Analysis

Dev. in Sedimental, 34, J. J. Fripiat (Ed.), Elsevier, New York, vi + 235 pp., 1982, \$46.50

Reviewed by Herman E. Roberson

Advanced Techniques for Clay Mineral Analysis is a collection of review articles dealing with nine analytical techniques: thermoanalytical methods, high resolution electron-microscopy studies, neutron scattering techniques, nuclear magnetic resonance, Mossbauer spectroscopy, electron spin resonance, ultraviolet and visible light spectroscopy, far infrared spectroscopy, and electron spectroscopy for chemical analysis (ESCA).

Each chapter includes some discussion of theory as it relates to a particular technique, but the main emphasis according to J. J. Fripiat, the editor, was to be directed toward summarizing recent developments of clay research applications. In fact there is an unevenness in this regard. Some authors (e.g., P. L. Hall on neutron scattering and J. P. Eberhart on high resolution electron-inicroscopy) have devoted a significant proportion of their articles to theoretical discussion, while other authors (T. J. Pinnavaia on elecand R. D. Mackenzie on thermoanalytical methods) have no general discussion of the-

Three areas of study reviewed have been extensively developed by chemists but have only recently received attention from clay researchers: Mossbauer spectroscopy, nuclear magnetic resonance (NMR), and electron spin resonance. Applications of Mossbauer spectroscopy in the study of clays are fairly extensive; as a result the author, B. A. Goodman, makes no attempt to present a comprehensive review. However, these topics which were selected (e.g., identification of oxidation states of iron and identification of iron-containing mineral phases at levels below their limits of detection by other more conventional techniques) will be of interest to many clay researchers. The review of NMR applied to water-clay systems is comprehensive; the clay

researcher attempting to get an overview in this research area may want to start here. Electron spin resonance applications which are discussed include studies of the orientation of hydrated metal ions on basal clay surfaces, mobility of interlayer ions, and a num-

ber of interlamellar metal complexes. In his review of high resolution electronmicroscopy application, J. P. Eberhart points out that this technique has already been apied with success in the study of several layer icates (primarily well-crystallized micas). The technique holds promise for the clay mineralogist who is willing to devote himself to gaining a thorough knowledge of the imaging process required for analysis.

ESCA, a relatively new technique, is one

that holds great promise for clay research. The technique, because of its high selectivity for the surface of the material being analyzed, has a range of applications that includes adsorption studies as well as studies re lated to the growth and alteration of clay

In general the articles are well written and achieve the primary goal: to acquaint the reader with some important analytical techniques which have been recently applied in studies of clay minerals. However, it should be pointed out that excellent review articles on many of the topics covered in this book have been published recently. This, along with the price (\$46.50), may curtail sales.

Herman E. Roberson is with the Dehartment of Geological Sciences, State University of New York at Binghamton, Binghamton, New York.

Climatic Geomorphology

. Büdel (Engl. Transl.), Princeton University Press, Princeton, N.J., xix + 443 pp., 1982, hardbound, \$50; cloth, \$18.50.

Reviewed by A. T. Grove

Although this book opens with a map of the world showing the continents divided into 10 morphoclimatic regions, this is not simply a text book on geomorphology organized dimatically. Rather it is a dissertation which at-

Books (cont. on p. 124)

ANNALES New 1983 **GEOPHYSICAE**

Editor-in-Chief: Stephan Mueller Institut für Goephysik ETH-Hönggerberg CH-8095 Zürloh (Switzerland)

As the official organ of the EUROPEAN GEOPHYSICAL SOCIETY, this newly created journal has been formed by incorporating the previously existing French
- Annales de Géophysique - and Italian

« Annali di Geofisica». It will accept original manuscripts written in English, but French, German, Italian or Spanish are permitted occasionally. No page charges

three fields:

SECTION 1: Physics of the Earth's and Planetary Interiors, Seismology and Tectonophysics Editor: R. MADARIAGA (Paris)

SECTION 2: Physics of the Hydrosphere and of the Earth's and Planetary Atmospheres Editor: R.R. VAN DER PLOEG (Stuttgart)

SECTION 5: Physics of the Interplanetary Medium, the Magnetosphere and the Upper Atmosphere Editor: H. REME (Toulouse)

Enhancement of resolution in selfadaptively deconvolved seismograms using short-time homomorphic wavelet estimation. M. KALKBRENNER, J.Y. KIM and J BEHRENS

Positive and negative ions in the stratosphere.

The spectral matrix, eigenvalues, and principal components in the analysis of multichannel geophysical data. J.C. SAMSON

Emission of myriametric radiation by coalescence of upper hybrid waves with low-frequency waves. K. RONNMARK

Observations of large scale F-region

at 7774 A and 6300 A.

Y. SAHAI, J.A. BITTENCOURT,

N.R. TEIXEIRA, H TAKAHASHI

irregularities using airglow emissions

Modelling of stratospheric ions: a first G. BRASSEUR, A. CHATEL

Annuel subscription (6 issues): 600 FF Foreign countries - 430 FF France.

Special subscription rate for individual members of EGS.

CDR-Centrale des Revues. 11, rue Gossin 92543 Montrouge Cedex-France

@ gauthier-villars

New and Recent Titles in Geophysics from Academic Press...

ICARUS

International Journal of Solar System Studies Editor: JOSEPH A. BURNS

Acting Editor: JOSEPH VEVERKA A repository for the major papers on the planetary sciences, Icarus publishes original contributions to the field of solar system studies. The papers report the results of current observational, experimental, and theoretical research on the astronomy, geology, meteorology, physics, chemistry, and biology. The journal is published in affiliation with the Division for Planetary Sciences of the Amor-

ican Astronomical Society. Volumes 53-56 1983, 12 issues and Cumulative Subject Index to Vols. I-52. Annual Subscription Rate, U.S.A. and Canada: \$384.00 Outside U.S.A. and Canada: \$434.50 ISSN: 0019-1035

LITHOSTRATIGRAPHIC ANALYSIS OF SEDIMENTARY BASINS

C. E. B. CONYBEARE "The book is a well arranged and comprehensive review of concepts and methods, spanning the evolution and development of basins, depositional environments, analysis and presentation techniques, and practical applications. . . . liberally illustrated with examples drawn from many parts of the world. . . . geologists and geophysicists working on the periphery of the specialist area would be well served by personal copies."

—David King in SEARCH

1979, 555 pp., \$55.50 ISBN: 0-12-186050-7

Journal subscriptions are for the calendar year and are payable in advance, Prices include postage, handling, and air freight where applicable. Individuals may request information regarding personal subscription rates, Please direct all orders and inquiries to the Journal Subscription Fulfillment Department

On book orders, send payment with order and save postage and handling. All prices are in U.S. dollars and are subject to change

ENERGY AT THE SURFACE OF THE EARTH An Introduction to the

Energetics of Ecosystems DAVID H. MILLER A Volume in the INTERNATIONAL GEOPHYSICS Series

"All the topics are here, from the general to the specific, physical through biotic: direct solar radiation, resultants of the upward and downward radiation fluxes, fixing of carbon by ecosystems, and the release of carbon fixed in ecosystems. . [Miller] provides data where data are appropriate, and offers personal discussions and interpreta-tions."—G. M. Goodwell in AMERICAN SCIENTIST

1981, 534 pp., \$27.50 ISBN: 0-12-497152-0 (Softcover) 1981, 534 pp., \$54.50 ISBN: 0-12-497150-4 (Hardcover)

To obtain examination copies of Energy at the Surface of the Earth for class use, address your request to the attention of the Sales Department, and give the title and estimated enrollment of your course.

SOLITONS AND NONLINEAR **WAVE EQUATIONS** R. K. DODD, J. C. EILBECK,

J. D. GIBBON, and H. C. MORRIS This volume describes the fundamentals of solitons and emphasizes the theory and applications of the classical soliton. It covers the study of the solution of certain equations solvable by the inverse scattering (spectral) transform and the investigation of classes of physical systems that give rise to these equations. Areas of physical application considered include plasmas, classical fluids, and

1982, 640 pp., \$45.00 ISBN: 0-12-219120-x

quantum optics.

OCEANOGRAPHY VOLUME 8 Edited by J. P. RILEY

and R. CHESTER The newest volume in this authoritative sories presents the latest advances in chemical oceanography from the per-spectives of the chemist, the biologist, and the geoscientist. Topics covered in-clude the effect of pressure on reactions in the sea, trace metal distribution, hydrothermal fluxes, interstitial waters, marine photochemistry, and the GEO-SECS program. The chapters are liberally illustrated with diagrams, tables, and photographs. Wherever possible, a mathematical approach to the subject has been adopted.

1983, in preparation ISBN: 0-12-588608-x

MOUNTAIN BUILDING **PROCESSES**

KENNETH J. HSÜ This work is based on the proceedings o the Symposium on Mountain Building held at the Geological Institute, ETH Zurich, 14th-18th July 1981. Generously illustrated, the papers in this volume dis cuss all aspects of the various plate in-teractions, with particular emphasis on the collision type of orogeny. The first section takes a theoretical and phenome nological approach and covers the tectonic processes associated with mountain building. In the second section historical case studies of several prominent orogenic belts illustrate the operation of auch processes. The authors are interna-tionally recognized and include former presidents of the Geological Societies of America and London.

1983, 288 pp., 8¾" x 11¾", \$72.50 ISBN: 0-12-357980-5

ACADEMIC PRESS, INC.

A Subsidiary of Harcourt Brace Jovanovich, Publishers New York • London • Toronto • Sydney • San Francisco 111 FIFTH AVENUE, NEW YORK, N.Y. 10003

ADVANCES IN GEOPHYSICS VOLUME 25 THEORY OF CLIMATE

Edited by BARRY SALTZMAN CONTENTS: HISTORY AND APPLICATION OF GENERAL CIRCULATION MODELS/J. Smagorinsky, The Beginnings of Numerical Weather Prediction and General Circulation Modeling: Early Recollections. S. Manabe, Carbon Monoxide and Climatic Change. STATISTICAL/DYNAMICAL MODELS/G. S. Golitana. Almost Empirical ELS/G. S. Golitsyn, Almost Empirical Approaches to the Problem of Climate, Its Variations and Fluctuations. G. J. Shutts, Parameterization of Traveling Weather Systems in a Simple Model of Large-Scale Atmospheric Flow. B. Saltz-man, Climatic Systems Analysis. RADI-ATIVE, SURFICIAL, AND DYNAMI-CAL PROPERTIES OF THE EARTH-ATMOSPHERE SYSTEM/C. Ohring and A. Gruber, Satellite Radiation Observations and Climate Theory. R. Dickinson, Land Surface Processes and Climate—Surface Albedos and Energy Balort and J.P. bal Angular Momentum and Energy Balance Requirements from Observations. 1983, 512 pp., \$64.00 ISBN: 0-12-018825-2

AND GLOBAL TECTONIC **SETTINGS** A. H. G. MITCHELI

and M. S. GARSON

The formulation of the global tectonic hypothesis has led to the evolution of a new understanding of the genesis and setting of ore deposits. The book identifies the major settings at plate boundaries and within plates from the geological record of the next 2500 millennia. logical record of the past 2500 millennia The authors emphasize the treatment of rocks as ores as part of the stratigraphic succession or intrusive body in which they are formed.

1982, 422 pp., \$51.00 ISBN: 0-12-499050-9

313212

123

MINERAL DEPOSITS

A Volume in the ACADEMIC PRESS GEOLOGY Series

Books (cont. from p. 123)

tempts to provide an explanation for the morphology of the earth. L. C. King had the same ambition, as had W. Penck and, more diffidently, W. M. Davis. One may continue to be impressed by the clarity and visual attractiveness of Davis's exposition, though dissatisfied with the integration of its elements; W. Penck's ideas, notably his mechanism for piedmont stairway development, are more difficult to appreciate; King's explanation of upland plains in terms of scarp retreat over immense distances as the result of uplift generated by the disruption of the continents fails to convince most geomorphologists.

A basic problem to be solved is the means by which the land surface is lowered to produce extensive plains developed on solid rock, which may then be dissected, with the preservation of extensive fragments apparendy little modified.

The German text from which this book is derived appeared in 1977. Budel was a pupil of Bruckner and Albert Penck. A splendid picture of both of them precedes the contents, one holding an umbrella, the other a hammer, both with pocket watches and seated on a pile of logs. These details are possibly significant in the light of what follows. Of the references, which number over 850, nearly three quarters are in German; 41 are to Budel himself, the earliest dating from 1933; many are from German university journals with papers by Bremer, Louis, Mensching, and Troll particularly well represented. H. F. Garner's "The Origin of Landscapes," proba-

bly the closest approach in English to Büdel's ters and the production of abrasive trost debook, is unfortunately attributed first to Carner and then to Gardner.

The gist of Büdel's argument is that landscapes can be explained in terms of the following sequence of events and shifts in cli-

(1) In Tertiary times, seasonally wet warm

climates extended from the equator to the poles and etchplains were created worldwide by the mechanism of double planation. This is assumed to result from the fact that chemical weathering at the base of the weathered layer is more intense than anywhere else in the world because the soil fauna and flora supply large quantities of carbonic acid, and the water is renewed every rainy season. Along joint planes decomposition takes place particularly deeply giving great thicknesses of grus with core stones. Rock masses protruding above the soil are regarded as exempt from such weathering and between these shield inselbergs, 'Surface wash in the shifting network of tiny rainy season rivulets and wash channels' is regarded as 'the decisive process causing planation' (p. 146).
(2) By the Late Pliocene, climatic zones

were becoming differentiated with the developinent of polar icecaps, and etcliplain development was increasingly confined to middle and low latitudes. 'The main areas of today's rivers became fixed at medium elevations in Central Europe' (p. 259), and high valley sys-tems formed in the Alps give the trough

sponding with the direction of the trade (4) The Holocene period has been so short that only a few percent of the relief can be ascribed to it. This is particularly the case in the middle latitudes, best known to most genmorphologists, where processes are peculiarly weak except where they have been artificially accelerated by deforestation.

(3) The Pleistocene is viewed as involving

in its last quarter a four told expansion and

retreat of icecaps. Funding processes in mul-

die latitudes resulted in the rapid mersion of

box valleys 100 to 300 in deep, largely be-

cause of the disruptive effects of her finds' ground ice immediately below the active lay-

er, which 'promotes both vertical and lateral

erosion and is the driving torce causing the

periglacial frost debris zone to be morpho-

ting (p. 105). At the same time, inselliergs

were removed by frost action. In warm de-

sheets, and basal surfaces of etcholaris were

extensively revealed by stripping away of the weathered layer, especially along lines corre-

serts, soil sheets were replaced by rubble

lynamically the zone of excessive valley out-

This scheme is exemplified by various 1cgional examples taken mainly from southern Germany and southeast Europe.

Budel's exposition can be criticized on various grounds. Too often he assetts without presenting evidence. More support is needed, for instance, for the claim on p. 124 that the intense chemical weathering taking place in the peritropical zone 'exceeds all chemical

weathering found in any other climate by really self-evident, therebue, that there is cry of the beamodical some must carl conmons amounts of dissolved mater, for this the tropics above all that many torks are par tropage are buppe to solution, (b. 150). In our to sust proper is for little attempt to po-vide eliconological control through the used existing isotopic dates of intrusive and estinsive tooks. Nevertheless one can welcome the to thright emphasis on the reliance of precut-gray to this on brockess no jouker obust ing on them and the acute commens bad on held observation. I particularly appreciaed a section on the burial of Olympia (pp.

Perhaps this volume, by teintroducing Bir. ish and American geomorphologists to a fadirion they abandoned a generation ago, sit lead some of them to attempt a reassessment of the morphology of the earth. While it could well be comparable to that of Buddy such a reassessment should take fuller account of the advances made over the last to decades in knowledge of plate tectoric, the Cenozoic climatic record, and geomorphic processes, and it should not neglect the enprinced abbrearition remote sensing has be en us of the form of the surface and whalis beneath ir.

A. T. Grove is the director of the African State Center, University of Cambridge, Cambridge, United Kingdom

Classified

RATES PER LINE

Positions Wanted: first insertion \$1.75, additional inserdons \$1.50. Positions Available, Services, Supplies, Courses,

and Announcements: hast insection \$3.50, additional insertions \$2.75.

Student Opportunities: host insertion free, additional insertions \$1.50.

There are no discounts or commissions on classified ads. Any type style that is not publisher's choice is charged at general advertising rates Loris published weekly on Tuesday. Ads must be received in writing on Monday of the week prior to publication

Replies to aids with box numbers should be addressed to Box ... American Geophysical Union, 2000 Florida Avenue, N.W., Washington,

For turther information, call toll-free 800-424-2488 or, in the Washington, D. C., area.

POSITIONS AVAILABLE

POSITIONS AVAILABLE.

Postdoctoral Position in Physical Oceanography. A postdoctoral appointment in physical (xeanography will be available beginning September, 1985 in the College of Marine Studies, University of Delaware, Newark, Delaware. The initial appointment will be for one year with probable extension for a second year. The salary will be \$20,000-\$24,000 per year, depending on experience. Funds for the position will be available largely from a grant by NSF for conduct and analysis of a field observational study of the shelfbreak front in the Middle Atlantic Bight.

The person obtaining the appointment would be responsible for a portion of the planning and execution of the field study, much of the subsequent data analysis and interpretation, and teaching of one graduate level course in physical oceanography each year. The successful applicant must have received the Ph.D. in physical oceanography or a closely related field by the starting date of his appointment. Preference will be given to applicants with direct experience in field observations.

To apply send a complete resume and the names of three references to Professor R. W. Carvine, College of Marine Studies, University of Delaware, Newark, DE 19711. (Telephone: (302) 738-2169).

The University of Delaware is an equal opportunity/affirmative action employer.

Happier You'll Be With You Want Results Advertising

EOS' weekly frequency means rapid publication and delivery of your message.

Communications in EOS result in accelerated sales and recruiting . . . plus a higher level of awareness of your product or service.

If you are impatient for results, weekly exposure in EOS is the

Call Robin Little Toll free 800-124-2488 She thinks impatience is a virtue.

Postdoctoral Research Associate in Geophysical Fluid Dynamical University of Southern California. Applications are invited for a research appointment in Geophysical Fluid Dynamics. The successful applicate to successful applicate to successful applicate. appointment in Geophysical Fluid Dynamics. The successful applicant would be expected to spend the majority of his time working on laboratory models of coastal upwelling. Appointment can begin immediately. Applicants should possess a recent Ph.D. degree in a relevant area of engineering, occanography or physics. A complete resume and at least three letters of recommendation should be sent to Prof. T. Maxworthy, OHE 430, Department of Mechatical Engineering, University of Southern California, Lot Angeles, CA 90089–1453.

shoulders of glaciated valleys. Pediment for-

mation began as essentially a trimming proc-ess mainly confined to deserts with cold win-

Assistant Professor/University of Alberta. The Department of Physics at the University of Alberta invites applications for a tenure track position at the level of an Assistant Professor in Physics in any of

level of an Assistant Professor in Physics in any of the following areas:

1. Astrophysics and Astronomy;
2. Geophysics (Electromagnetic methods);
3. Theoretical Physics (Mediunt Energy, Particle Physics, Relativity and Cosmology).

The 1982/83 salary range for an Assistant Professor is \$27,720–839,820 per annum.

Applications will be received until May 1, 1983, and the expected appointment date is July 1, 1983. The Department of Physics offers both undergraduate and graduate degrees in Physics and Geophysics. The Department currently consists of 47 faculty Members, 36 Research Associates and Post-Doctoral Fellows and 50 Graduate Students.

Candidates interested in applying should submit a curriculum vitae plus the names of three (3) referees to:

Dr. A. N. Kamal

Dr. A. N. Kamal
Chairman
Department of Physics
University of Alberta
Edmonton, Alberta, Canada
T6G 211
The University of Alberta is an equal opportunity
employer but, in accordance with Canadian immigration requirements, priority will be given to Canadian citizens and permanent residents of Canada.

President/University Corporation for Atmospheric Research (UCAR), Boulder, Colorado. The Search Committee of the UCAR Board of Trustees invites nominations and applications for the President and Chief Executive Officer of UCAR. The Board of Trustees expects to fill the position promptly and seeks to lave the individual in office on July 1 or as soon after I July 1983 as possible. UCAR is a private, nonprofit consortium of 48 U.S. and two Canadian institutions with doctoral programs in the atmospheric sciences and related fields, UCAR's major activities consist of overseeing the programs of the National Center for Atmospheric Research (NCAR) in Boulder, Colorado, and the National Scientific Balloon Facility (NSBF) in Palestine, Texas, and managing their operations; the planning and conducts of 50 in Palesine, Texas, and managing their operations; the planning and conduct of special, cooperative atmospheric research programs; and foatering atmospheric research in the national interest.

To be considered for the UCAR presidency, a person should have demonstrated imaginative scientific and management leaderstell in the scientific and management lead iffic and management leadership ability of high quality and specific knowledge of the atmospheric and related sciences. A complete description of the duties and responsibilities of the President and qualifications for the candidates may be obtained by writing to Dr. Donahue (address below). Applications and nominations, which should be postmarked no later than 6 May 1983, should be addressed to:

Dr. Thomas M. Donahue
Chairman, Search Committee

UCAR
P.O. Box 3000

UCAR
P.O. Box 3000
Boulder, Colorado 80307
UCAR is an equal opportunity, affirmative action

Faculty Positions/The University of Iowa. The Department of Physics and Astronomy anticipates one or two openings for tenure-track assistant professors or any rank in August 1983. Preference will be given to experimentalists in any area for the tenure-track positions. Current research interests include astronomy, atomic, condensed matter, elementary patricle, laser, nuclear, plasma, and space physics. The positions involve undergraduate and graduate teaching guidance of research students, and personal research. Interesteen persons should send a resume and a statement of reasons should send a resume and a statement of reasons interests, and have three letters of recommendation sent to Search Committee, Department of Physics and Astronomy, The University of Iowa, Iowa The University of Iowa is an excel-The University of Iowa is an equal opportunity/

Chairman—Department of Geological Sciences, Wright State University. The Department of Geological Sciences, invites applications for the position of chairman, to be appointed September 1984. We seek a dynamic individual with administrative talent and an appreciation for research and practice-telated educational activities. Rank is at the full professor level and no restrictions have been always to the contractions. ed educational activities. Rank is at the full profession level and no restrictions have been placed on arcas of specialization. The department is active with 12 faculty and an emphasis on professional practice, yet maintaining a firm communication basic research.

Send a letter of application, corriculum virae and Send a feuer of application, confliction vitae and names of three references to:

Chairman, Search Committee
Department of Geological Sciences
Wright State University
Davion, OH 45-45,
Wright State University is an affirmative action/could apportunity considerer. Classing data for the

al opportunity employer. Glosing date for the ition is October 31, 1983.

Sedimentary Geology/University of Pittsburgh. The Department of Geology and Phanetary Science invites applications for a tenure-stream faculty position in sedimentary geology at the assistant professor level, which is expected to be upon beginning September 1983. We seek randidates with expertise in one or more of the following fields: sedimentary petrology, sedimentation, stras: sedimentary petrology, sedimentation, stra-aphy, sedimentary geochemistry, and economic

geology of non-metallic deposits. The position is volves development of a vigorous research progra-supervision of graduate student research, and teab-ture of source. ing of undergraduate and graduate courses.

Applicants should submit a letter of application. statement of research interests, transcripts, via, and

Dr. E. G. Ludiak, Chairman, Department of Go-ogy and Planetary Science, University of Finburg Pittsburgh, PA 15260

Postdoctoral Position in Solar-Terrestrial Physics. The Center for Atmospheric and Spac Sciences of Utah State University has a postdoctor position in solar retrestrial physics. The position for one year but has renewable opportunities the textipient is expected to develop theoretical model to describe the large-scale flow of plasma in the restricted impositions and ones. to describe the large-scale flow of plasma in twee restrial ionosphere and magnetosphere and to subthe stability of the flow. Candidates should have Ph.D. and a background in space physics are plasma physics. Send letter of application, resurtant the names and addresses of two references to April 30 to Professor R. W. Schunk, Center for Amostuberic and State Sciences, Unit State University. mospheric and Space Sciences, Utah State University, Logan, Utah 81322 (801-780-2974).

Utah State University is an Albumative Adion

DEAN OF GEOSCIENCES **TEXAS A&M UNIVERSITY**

Texas A&M University Invites applications and nominations for the position of Dean of Geosciences

QUALIFICATIONS: Dean of the College of Geosciences should have national or international reputation for excellence in teaching, research, and/or administration in at least one of the fields of geoscience represented at Texas A&M University. (Geography, Geology, Geophysics Meteorology, and Oceanography).

Evidence of such competence should be demonstrated by the individual's history relative to publications, honors, professional associations, participation in national councils and advisory panels and in positions and appointments. This history should include evidence of appreciation and understanding of management problems on all levels demonstrated ability to make decisions to administer programs, delegate authority and establish policy with sensitivity, diplomacy, fairness, firm-

Of much importance for the College of Geosciences at Texas A&M University is the Dean's ability to establish favorable rapport with industry and government, particularly with reference to sources of research funding. The Dean should be characterized by his/her ability to communicate effectively. The Dean must be endowed with sufficient desire and vision to lead the College to a high position of national promi-

TEXAS A&M UNIVERSITY: Texas A&M University is a rapidly-growing four-year, state-supported university with enrollment exceeding 35,000 students. The university is located in College Station, TX approximately 90 miles from Houston, TX. The twin cities of Bryan and College Station have a combined population exceeding 100,000 people.


CONDITIONS OF EMPLOYMENT: This is a 12-month position available

on or before September 1, 1983. Salary is competitive and negotiable. APPLICATION PROCEDURE: The application should have at least three references who are familiar with the applicant's or nominee's background in higher education administration.

The deadline for application is April 15, 1983, Send letters of application and nomination to:

Dr. Charles E. McCandless Associate Vice President for Academic Affairs and Chairman, Search Committee Texas A&M University College Station, TX 77843.

Texas A&M is an equal opportunity employer.



Applications are invited for appointment to the above post with effect from 1 February

1984.
The University wishes to appoint a scientist with an established research record to take responsibility for the further development of research and teaching in his recently created department. Candidates currently accupying either academic or non-academic positions should have a wide knowledge of modern instrumental analysis and his requirement will be of prime importance in making the selection for the post. Appointment, depending on qualifications and experience, will be made in the salary range R23 109 to R30 255 per annum, to which is added an annual service bonus of nearly one month's salary.

Applicants should submit a full resumé and the names and addresses of three referees whom the University may approach. Further Information should be obtained to the Registrar (Attention: Appointments Office), University of Cape Town Private Bag, Rondebosch, 7700, South Africo, Office), University 17700, South Affice, by whom applications (qualing ref. no. BM/332) must be received not later than 24 June 1983.

Faculty Position/Physical Oceanography, University of Maryland Center for Environmental and Estuarine Studies (UMCEES). The Horn Point Environmental Laboratories of UMCEES invite applications for a tenure-track (rank open) position for a physical oceanographer. The physical oceanographic program is young and developing, so that candidates with a wide range of research interests will be considered. Opportunities exist for oceanographers with experimental, theoretical, or numerical modeling skills. In addition, the strong chemical and biological programs provide opportunities for interdisciplinary research. Although some preference will be given to applicants with interests in continental shelf and estuarine circulation processes, the primary criterion for selection of the successful candidate is the ability to develop a strong research program. The closing date for applications is May 16, 1983. Applications and a list of references should be sent to:

Dr. William C. Bolcourt
Chairman, Search Committee
University of Maryland
Center for Environmental and
Estuarine Studies
Horn Point Environmental I abstrato miental Laboratories

P. O. Box 775
Cambridge, Maryland 21613.
The University of Maryland is an Affirmative Ac-


Hydrogeologist. Applications are invited for a starting in January 1984. Rank and salary are de-pendent on qualifications and experience. Ph.D. de-gree and registration as a professional engineer de-sirable.

Responsibilities will include undergraduate and graduate teaching and graduate research in hydrogeology and engineering geology. Applicants should have experience in field groundwater investigations of periods. have experience in neid groundwater investigations of regional site specific nature/groundwater re-source analyses/waste disposat/groundwater contami-nation studies. Experience in applying remote sens-ing to hydrogeology and engineering studies would be before

be helpful.

Send resume and names of three references to J.
J. Finney, Head, Geology Department; Colorado
School of Mines, Colden, Colorado 80401. Applications will be accepted until September 15, 1989.

The Colorado School of Mines is an affirmative action/equal conocupity employer.



The successfull applicant will conduct research primarity in the field of Economic Geology and will also do some teaching within the Department. He or she will work closely with Professor L Minter in establishing a research programme in Economic Geology at the University of Cape Town.

Applications any or Cape I own.
Applications and the applications and experience in the solary range R12 657 to R22 173 per annum. Staff benefits include an annual bonus of nearly one month's solary, pension, medical ald and a housing subsidy subject to Stafe regulations.
Applicants should subside a surfacilities and the solary, pension, medical ald and a housing subsidy subject to Stafe regulations. Applicants should submit a curriculum vide storing research interests, age, present solarly, experience and qualifications, the date day could be assumed and the names of three referees.

Further information is obtainable from Professor A M Reid, Department of Geology, University of Cape Town, Private Bag, Rondebosch, 7700, South Africa, by Whom opplications should be received by 31 May 1983.

The University's policy is not to discriminate on the grounds of sex race or religion.

Geochemiat. A career position is available in the Earth Sciences Department at Lawrence Livermore National Laboratory, Livermore, California, for a Geochemist, or a person with related training or experience, to provide geochemical modeling support to the evaluation of repository sites for the disposal of nuclear waste.

port to the evaluation of repository sites for the dis-posal of nuclear waste. The work involves developing and extending ex-isting geochemical computer codes, updating ther-modynamic data buses and experimental work in support of and to validate the geochemical models. Requires knowledge of FORTRAN, an advanced degree or equivalent in Geochemistry, Physical Chemistry or Inorganic Chemistry, and experience in developing and using computer codes to solve geochemical problems.

geoclemical problems.

Send detailed resume with full employment history and objectives to Art Wong, Professional Employment Division, LAWRENCE LIVERMORE NATIONAL LABORATORY, P. O. Box 5510, KES-33, Livermore, CA 94550. U.S. citizenship is

in equal opportunity employer M/F/H/V.

Mineralogy and Petroleum Geology/Eastern Washington University. The Department of Geology is seeking applicants for a faculty position in mineralogy and one in petroleum geology. Doctoral degree required, Mineralogist to teach mineralogy and averay differaction/fluorescence. Petroleum geologist with industrial experience to teach petroleum geology and some area of soft rock geology. Both positions require teaching undergraduate/graduate courses in area of specialization plus introductory geology. Development of strong research program expected. Positions start September 1, 1983. Evaluation of application, vita, and 4 letters of recommendation to Dr. Eugene Kiver, Search Committee, Department of Geology, Eastern Washington University, Cheney, Washington 99004.

Research Position/Space Physics. The Space Physics and Astronomy Department at Rice University seeks applicants for one or more full-time research positions within the department. Successful applicants) will play key role(s) in the development of theoretical three-dimensional models of the Earth's electromagnetic field. Applicants should have knowledge of, and interest in, at least one of the following areas: solar-wind magnetosphere interactions, magnetosphere-ionosphere coupling, ionosphere-atmosphere coupling, atmosphere clectricity. Experience and/or interest in numerical modeling is an important consideration.

Title and salary level commensurate with experience, ranging from one-year Research Associateship trenewable in subsequent years depending on performance to open-ended Research scientist appoinment in the Center to Space Physics. Please send resume and names of three professional references to T. W. Hill or R. A. Wolf, Space Physics and Astronomy Department, Rice University, Houston, TX 77251.

The University is an equal opportunity/affirmative

The University is an equal opportunity/affirmative

Staff Scientists/Systems Analysts. Research and Data Systems, Inc. has openings available for Stali Scientists, Systems Analysis and Programmer/Analysts to work in areas involved in the processing and application of data from satelline based remote sensing systems. Particular needs involve the analysis and processing of Earth Radiation Budget, Microwave, AVHRR and LANDSAT data. Needs also exist in the areas of interactive image graphics, software engineering, realtine processing and satelline data communications. Successful candidates will have an advanced degree in meteorology, physics, engineering, mathematics, or computer science. Hatchware background doubt include 1534, 1447. CYBER or 149-1000 equipment. Send resume in confidence to: lysta to work in areas involved in th

ce to: Research and Data Systems, Inc. 10300 Greenbeh Road, Suite 206 Lanham, Maryland 20706 Telephone: (301) 390-6100.

SERVICES, SUPPLIES, COURSES, AND ANNOUNCEMENTS

May 9-13, 1983. "Planning, Design, and Operation of Real-Time Data Monitoring Systems and Applications." \$750.00. Topics include remote-site systems are applications." cations." \$750.00. Topics include remote-site sensors and telemetry equipment, communications systems options (satellite, radio line-of-sight, telephone, etc.), data collection platforms, computer systems software, data base design and management, applications that incude flood control, irrigation, hydropower production, hydrofogical and meteorological data collection, water quality, air pollution, acid rain, groundwater levels and quality, etc., economics of real-time data collection, use of radar information, current status of hardware, computer graphics and computer imagery for remote sensing-for additional information, contact Dr. Raul S. McQuivey, Sutron Corporation, 11150 Main Street, Fairfax, Virginia (703) 591-8010.

Engineering Analysis of Fluvial Systems

August 1-4, 1983 Copper Mountain Resort, Colorado

State-of-the-art design techniques oriented to solving practical engineering problems associated with watersheds and rivers are presented in this four day short course. The course is designed to enhance the participants' insight mathematical modeling techniques. Design applications are illustrated with numerous case studies. The recently published hardbound text Engineering Analysis of Fluvial Systems will be provided to each participant.

Lecturers: D.B. Simons, R.M. Li, and P.F. Lagasse

Registration fee: \$450 U.S. Dollars

Contect: Dr. Peter F. Lagasse Simons, Li & Associates, Inc. P.O. Box 1816 Fort Collins, CO 80522 (303) 223-4100

STUDENT OPPORTUNITIES

IBM Environmental Engineering Graduate
Fellowship. Colorado State University invites applications from atudents interested in a Ph.D. engineering program with specialities in water quality hydrology and water quality monitoring. The fellowship provides a \$1,000/month stipend, all tuition and fees, books, research expenditures and travel funds. For further information or application,

Dr. Robert G. Ward
Department of Agricultural and
Chemical Engineering
Colorado State University
Fort Collins, CO 80525

RESEARCH POSITION IN GEODYNAMICS-ASTROMETRY

At the Lunar Satellite Ranging Facility Haleakala Observatory, Maui, Hawaii

The University of Hawaii's Institute for Astronomy has an immediate opening for a Researcher.

The selected applicant will work closely with the Observatory Project Manager to ensure an effective program of ranging on artificial satellites and the moon; will also conduct a research program in an area of interest to the facility; and will be a resident on the Island of Maui. Minimum Qualifications: Ph.D. in Geosciences, Astronomy, or Physics and a proven record as a researcher as demonstrated by a list of publications and recommendations of peers.

The position is full-time, federally funded, subject to annual contract renewal, and continuation dependent upon availability of funds. Salary will be commensurate with qualifications and experience.

Interested individuals should send a bio-bibliographical summary together with the names of two or three people who have knowledge of the applicant's professional abilities to: Ms. Carol Yoshida, Personnel Officer, Institute for Astronomy, 2680 Woodlawn Drive, Honolulu, Hawaii 96822. Further details can be obtained from Dr. John T. Jefferies, Director at (808) 948-8566. Applications should be postmarked no later than April 29, 1983.

AN EQUAL OPPORTUNITY/AFFIRMATIVE ACTION EMPLOYER

Membership Applications Received

Applications for membership have been recived from the following individuals. The letter after the name denotes the proposed primary section affiliation; the letter A denotes the Atmospheric Sciences section, which was formerly the Meteorology section.

Regular Member

David R. Bazard (GP), Stephen Blake (V), Soman Chacko (S), Don W. Chorley (H), Edward Cranswick (S), David A. Dinter (T), Roper R. Faton (H). Fred G. Everdale (O). Ingrid P. Fotino (O), Christopher Fox (T), Robert H. Gilkeson (H), Douglas D. Given (S), Arthur Goldstein (T), Edward J. Green (O), Margaret Hellweg (S), Alexander D. Jack (S), Cathy J. Janik (V), Douglas W. Johnson (A), Jacob Kushnir (O).

Robert C. Livingston (SA), Asger Lundbak (SS), Mario J. Martinez (H), H. Lawrence McKague, David McKirdy (GP), George H. Mount (SA), Stanley K. Nazalewicz (G), S. L. Passman (T), Kristine Rock (SS), Ralph Rogers (T), Richard S. Scalan (O), Roberto Scandone (V), Daniel S. Spicer (SM), Kenneth R. Stalder (SM), Donald H. Stedman (A), Bryan Tapp (T), Murad Taqqu (H), Janet K.
Thompson (O), John J. Ward (H), Randall A.
White (S), Derek A. Widmayer (H), Douglas

Student Member

Yemane Asmerom (V), Donna K. Blackman (T), Leslie J. Blythe (H), Jeffrey A. Brooks (P), Grover S. Buhr (S), Yildirim Di-lek (T), David G. Evans (S), Guy Gelfenbaum (O). Kirsti Granlund, Nancy W. Hinman (O).

AGU GIFT Recognition for Your Support

> Supporting Member \$80.00/year

agu

GIFT

List appears in Eas. Special recognition national meetings.

Life Supporting Member \$1,500 List appears in Eos. No further dues ob-

Sustaining Member \$5,000 Benefactor \$10,000

A certificate of appreciation from the president will be given for contributions totaling \$500 for the 5 years (1981-1985). Contributors of \$1000 or more will be individually recognized in a permanent display at the headquarters of the Union.

Hanaly M. Holail (V), C. D. Hull (V), Bruce Idleman (T), David A. L. Johnson (S), Steven Knapp, Kevin M. Konstanty (S), Robert P. Labelle (O), Nathaniel A. Litton (T), Robert S. Linzell (O).

Pagu D. Maniar (V) Patrick I. Neale (O) Jong-Sim Park (V), Sam-Keun Park, Michael Pound (S). Thomas T. Ruubel (E). Cymbia. T. Schramm (O), Chris Sherwood (O), Peifu I. Shih (SA), Bunny Sterin (H), Robert C. Symberlist (T), Todd M. Thornburg (O), Norbert Wild (SM), Dean Witte (S).

Associate Member

David N. Jenkins (H), Charles E. Mongan, Helen Musiafa (O), Kenneth O. Sizemore (SS), Eliza I. Woitaszek (T).

The June Bacon-Bercey Scholarship in **Atmospheric Sciences for Women** 1983-1984

Expressly for women intending to make a career in the atmospheric sciences. This monetary assistance, provided through a gift from June Bacon-Bercey, a noted meteorologist, will be given to a woman who shows academic achievement and promise. To qualify, candidates must be one of the following:

• a first-year graduate student in an advanced degree program in atmospheric sciences; an undergraduate in a bachelors

degree program in atmospheric sciences who has been accepted for graduate study; • a student of a 2-year institution of-

fering at least six semester hours of atmospheric sciences, who has been accepted for a bachelor's degree program, and who has completed all of the courses in atmospheric science offered at the 2-year institution.

Awardee selection will be made by the AGU Subcommittee on Women in Geophysics in consultation with the AGU Atmospheric Sciences Section.

For application forms contact: American Geophysical Union Member Programs Division 2000 Florida Avenue, N.W. Washington, D.C. 20009

462-6903 800-424-2488 outside the Washington, D.C. area Application Deadline May 1, 1983

. . .

- ----

Meetinas

Announcements

Call for Papers: Chapman Conference on Magnetic Reconnection

A Chapman Conference on 'Magnetic Reconnection' will be held at the Los Alamos National Laboratory, Los Alamos, New Mexico, October 3-7, 1983. Since its conception over three decades ago as a possible solar flare mechanism, magnetic reconnection has become a matter of substantial interest and probable importance not only for flares but also for other domains of plasma physics, including planetary and stellar magnetospheres and laboratory fusion research. The main emphasis of this meeting will be on magnetic reconnection in the earth's magnetosphere, but the perceived role of reconnection in other cosmic objects and in laboratory plasmas will also be treated to provide a full picture of

the present understanding of this process.

Specific areas of discussion will include the following topics: theories of reconnection and its anticipated signatures, reconnection at the earth's magnetopause, reconnection in the earth's magnetotail, reconnection in astronomical objects, reconnection in laboratory plasmas, computer modeling of reconnection, and directions for future research.

The 41/2-day conference will include morning sessions plus a mix of afternoon and evening sessions that will leave sufficient time for discussion. Poster papers will be encouraged and will be previewed and displayed so as to optimize their communication to Conference participants. Arrangements will be made to provide to attendees during the meeting a volume of extended summaries of the papers and to publish conference papers in book

All who are interested in attending and in receiving later information circulars should write to Magnetic Reconnection Meeting, AGU, 2000 Florida Avenue, N.W., Wash ton, DC 20009 (telephone, toll free: 800-424-2488 or, in the D.C. area, 462-6903).

To submit an abstract, follow the abstract format published in Eas, November 30, 1982. and January 18, 1983. There will be no abstract charge. All abstracts should be sent to Magnetic Reconnection Meeting, AGU, at the address in the previous paragraph.

ABSTRACTS DEADLINE JULY 1, 1983

Program Committee: E. W. Hones, Jr., Los Alamos National Laboratory; V. M. Vasy-liunas, Max-Planck-Institute for Aeronomy, Katlenburg-Lindau, FRG; F. Coroniti, Department of Physics, UCLA; D. N. Baker, Los Alamos National Laboratory; D. H. Fairfield, NASA Goddard Space Flight Center; A. Nishida, Institute of Space and Astronautical Science, Tokyo; J. Brackbill, Los Alamos National Laboratory; C. T. Russell, Institute of Geophysics and Planetary Physics, UCLA; B. U. O. Sonnerup, Radiophysics Laboratory, Dartmouth College, Student Travel: Limited funding is available

to support student travel expenses to the conference. To apply, write to AGU giving your educational background, your reasons for wanting to attend the conference, and your research interests. The awardees will be selected by AGU in conjunction with the program committee. Deadline for travel applications is July 1, 1983.

Hydrology Days Update

The AGU Front Range Branch Hydrology Days symposium will be held at the Colorado State University in Fort Collins, April 19-21. The 28 papers to be presented will cover

such topics as flow resistance of bombler-hed streams; snowmelt runoff simulation using the Martinec-Rango model; long-term models of phosphorous in completely mixed lakes; multilayered aquifer modeling; the rainfallrunoff process from a geomorphic-hydraulic perspective; infiltration, soil moisture redistribution, soil evaporation, and aquifer recharge in the HEC 1 (Hydrological Engineering Center) model; approaches for estimating regional snowpack equivalent; and variability of bed material transport and channel hydrau-lics with high washload concentrations.

The AGU Front Range Branch Hydrology Days (Eas, November 2, 1982, p. 838, and July 27, 1982, p. 589) is dedicated to Robert . Glover in recognition of his contributions to hydrology. Glover, on the faculty of Calorado State University from 1956 to 1963, is currently a faculty affiliate.

For more information, a complete program, and registration materials, contact [1.]. Morel-Seytoux, Department of Civil Engi-neering, Colorado State University, Fort Collins, CO 80523 (telephone: 303-491-5448).

Geophysical Year

New Listings

The complete Geophysical Year last appeared in the December 21, 1982, Eos. A boldface meeting title indicates spousorship or cosponsorship by AGU.

August 29-September 8, 1983 4th International Symposium on Water-Rock Interaction, MISASA, Japan. (Professor H. Sakai, Secretary-General, WRI-4, Institute for Thermal Services.) mal Spring Research, Okayama University, MISASA, Tottori-Ken 682–02, Japan.)

Water Resources Monograph 7

David F. Ribler, colors Urban Stormwater Hydrology

A practical quide to current methods & mode used in analyzing different types of stomoste management problems

Topics include interaction of land use and intestationater runoff . Stormwater planning in to inten metroples . Current design minial insued for water quality studies . Deskins methods for urban stormwater calculation than tunult processes. Pullution potential of tom-water. I tale of the transport system C Data of lection and instrumentation • and more

A major reference work for those involved in water terources research, graduate students, prac-ticing emphases and within planners

Contains illustrative step by step examples the principal large scale planning and designates runoff models, including such programs as STORM, SEMS TORM, ILLUDAS, SWMM, RUNQUAL, HSPF and others

 Blustrated • 280 pages • 30% Member discour List price \$18.00

Orders under \$50 must be prepaid American Geophysical Union 2000 Florida Ave. N W Washington, D C 20009 Call 800-424-2488 trill free • 462-6903 (local)

September 30-C)ctober 1, 1983 Pacific Northwest Regional Meeting, Bellingham Wash. (Myrl E. Beck, Jr., PNAGU, Department of Geology, Western Washington Sale College, Bellingham, WA 98225.)

Separates

To Order: The order number can be found at the end of each abstract; use all digits when ordering. Only papers with order numbers are available from AGU. Cost: \$3.50 for the first article and \$1.00 for each additional article in the same order. Payment must accompany order. Deposit accounts available.

Copies of English translations of articles from Russian translation journals are available either in unedited form at the time of their listing in EOS or in final printed form when a journal is published. The charge is \$2.00 per Russian page.

> Send your order to: American Geophysical Union 2000 Florida Avenue, N.W. Washington, D.C. 20009

Aeronomy

O410 Absorption and Scattering of Radiation
O200E DEMSITIES IN THE LONER MESOSPHERE MASSURED BY A
LINE SCANNING LATRAVIOLET SPECTRONETER
O. W. Rusch (Laboratory for Atmospheric and Space
Physics and Department of Atmospheric and Space
Physics and Department of Atmospheric and Space
Physics and Department of Atmospheric and Space
of Coloredo, Boulder, CO 80300), G.H. Mount, C.A.
Sanders, G.H. Lawrence and R.S. Eckman,
The Coane content of the earth's atmosphere between
into and 0.08 mb has been measured as a function of
latitude and season by an effraviolet spectrometer on
latitude and season by an effraviolet spectrometer on
mixing ratio is found to be highly variable in time
and space during the winter Ammisphere during January and
February at all pressure layers. The latitude
gradients mear spring sequence are relatively small, A
20°S, in January and February an 30°S is January and February. Geophys. Ses. Lett., Paper 310318

0440 Convection, diffusion, mixing, turbulence, and GAAO Convection, diffusion, mixing, turbulence, and fellout fellout fellout fellout fellout fellout fellout turbulence. And fellout turbulence fellout fellout turbulence fellout fell J. Grephys. Pds., Blue, Paper 340410

Electromagnetics

OID) Antennes
SECOLARCE CORES IN MON-MARKELLICH FLARMES-2
B. Thismann (Center for Atmospheric and Space
Sciences, Useh Static Datversity, Logan, Wich
Selection, Useh Static Datversity, Logan, Wich
Selection of Market State
Between Committee of the planes in described
by Earl stagings against last planes in described
by Earl stagings appoint item; a background population had an electron beau with variable beau
justance had an electron beau with variable beau
parameters. It is shown that the Corentor consvaluation had an electron beau with variable beau
parameters. It is shown that the Corentor consvaluation of the planes are a staging and selection of the lower harmed of
a showly may be a selection of the lower harmed of
a showly may recommise cons irraspect two at the

cagnitude of the blasting cagnetic field. When the bear whocity is sufficiently large with respect to the thermal velocity of the background electrons, a therecteristic wave pattern appears octained the resonance cones. It is shown that this pattern occurs because of a propagating wave for 9.5% (80.9%). When the bear density is low to 11), the upstressmenter come structure remains nearly unsaddled, and it can be used for plasma diagnostics to deduce electron respectature and density using theoretical results based on a Maxwellian plasma.

Rad. Sci., Paper 380296

Geochemistry

1410 Chemistry of the Atmosphere
EIRETICS AND RECEASES OF THE OXIDATION OF S(IV) BY
0200E IN AURODE SOLUTION WITH PARTICULAR REFERENCE
TO SO_CONVESTON IN ROWINDAY REPORTMENT COUNS
H. G. Manha (MASA Langlay Essearch Center, Hampton,
Virginia, 21665)
The kinetics of S(IV) exidation by ozone in equenum
solution have been studied at 10°C and 23°C in the pit
range 3.0 to 6.2 union stopped-flow spectrophotomeLey. The rate data are amplifically decreased by
I the rate data are amplifically decreased by

r = (k_L + k₂ |0^{pH})[s(tV)](03]

with $k_1 = 4.39 \times 10^{11} \text{ s}^{-4131/7}$ ($\pm 3 \times 10^4$) $\text{H}^{-1} \text{s}^{-1}$ and $k_2 =$ with kg = 4.39x10¹⁸ m^{-131/1} (±3x10⁴) H⁻¹s⁻¹ and kg = 2.56x10³ e^{-366/7} (±15) s⁻¹, at an ionic strangth of 0.01 N. Reaction Cate is unaffected by incident strong UV radiation, or the presence of Mo(11) or ys(I(1), but is unakly dependent on ionic strangth. The present tensits, in comparison with reported takes of voldation of S(IV) by hydrogan parceide, indicate that at typical pit lavels encountered in monorban tropospheric cloud water end at representative concentrations of 03 and H₂O₂ is the tropospheric considered by Oddard to the tropospheric considered in the property of the property of the property of the tropospheric considered in the concentrations of 0.3 and H₂O₂ is the tropospheric consideration by O₂ can be comparative with that by H₂O₂, or possibly even decising. The reaction appears to proceed by other than a fraction land reaction access, and a reaction school towards in 1s proposed. A rate expression consistent with this school is

r = (k_a + k_b[08"))(450])(03) where $k_{a} = 3.8 \times 10^{5} \text{ M}^{-1} \text{ and } k_{b} = 1.05 \times 10^{16} \text{ M}^{-2} \text{ s}^{-1}$ at 25°G. This empression correlates the present data in conjection with literature data over the pi range 1.0 to 6.2. (Suffice oridation, suiface formation, some cloud characteristics)

propa, cloud chamistry). Jacophys. Ses. Graco, Paper 3C0416 1416 Chemistry of the atmosphere
CHRACTERIZATION OF THE ATMOSPHERIC APROSOL OVER
THE PASTERN EJAMORTAL PACIFIC
THE PASTERN EJAMORTAL PACIFIC
THIS PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC
THIS PACIFIC PACIFIC PACIFIC PACIFIC
THIS PA

the enrichment being auhatumatically more pronounced (up to 50% or higher) for 1-4 µm diameter particles than for particles > 4 µm. Comparison of these data with as similar data set from samples collected over the Atlantic indicates that the departures from seasons composition are significantly larger for the Pacific. Differences in see-to-sir fractionation processes, probably involving binding of divalent tations to organic matter in the oceanic surface microlayer, are suggested as being responsible for these observations. (Acrosols, Scu sait, Pacific).

Pacific).

J. Geophys. Res., Gress, Paper 3COA17

1410 Chemistry of the Atmosphere

Irace Elements in Marine Aerosol Particles from
Enembels Atoli: 1. Concentrations, Sources, and
Temporal Variability
R.A. Buce (Center for Atmospheric Chemistry Studios,
Graduata School of Oceanography, University of Rhode
Island, Kingston, Rhode Island, O2881), R. Arimoto,
B.J. Ray, C.K. Unni and P.J. Harder
The concentrations of twenty-nine elements were
measured in aerosol particles collected in 1979 during
SEAREX experiences at Enemats Ad 2011 (1104), 1820[)
In the tropical North Pacific. The concentrations of
Ma, Mg, Cl. K. Ca, and Br were dominated by marine
sources; these elements had similar mass-size
distributions, and their atmospheric concentration
ratios (normalized to Ma) were similar to the
corresponding ratios in bulk seswater. Atmospheric
inputs of alaminos Ilicate particles from crustal
weathering controlled the aerosol particle
concentrations of Al, Sc, Ma, Fe, Co, Cs, Ba, Co, Eu,
HH, Ta, and Ih. Mean concentrations of these
crustally derived elements decreased by an average of
91 percent (± 4.1 percent) from the local dry season
(April to Nay) to the wet season (July to Augusti;
this general decrease was attributed to the abalamoni
of dust storms in Asia. At those the influx of dust
from Asia dominated the concentrations of Y, Cr, Rb,
and Cu in aerosol particles, but when dust
concentrations decreased, non-crustal sources for
these elements became apparent: A fourth group of
elements (Zn, Se, Ag, Cd, Sb, I, and Pb) exhibited
average atmospheric concentrations which were higher
than those expected from the flux of sea salt or the
dispersel of mineral aerosol particles. Enrichments
of these trace elements relative to average crustal
material increased as the atmospheric dust
concentrations subsided. (Marine aerosol particles,
trace alsomets).
J. Geoghys. Res., Greeg, Paper 3COA15

Hydrology

J125 Glaciology
On THE OYNAMICS OF THE ICE SHEETS II
P. Halfar (Max-Planck-Institut for Neteorologie, Hamburg, FRC)
The equation which describes the motion of an Ice sheet under its own weight has a cylindrically symmetric similarity solution. The initial value problem of the equation of motion which has been linearized in the coulations from this shallarity solution in terms of significant of the solution in terms of significant of the solution in equation. The linearized equation detarmines for all ice sheets of the same volume an asymptotically stable solion. The results are compared with the corresponding results of the J. Geophys. Res., Green, Paper 3COMO. J. Gaophys. Res., Green, Paper 300340

Oceanography

4713 Circulation
SOUTHWARD SUBSTRIVACE FLOW BELOW THE SCHALL CURRENT
D. R. Quadianel (Institut fift Mearaskonde, University,
Bambura, 2000 hasburg 13, F.2.g.), F. Schott (University)
of Miani, Florida 33149).
The existence of a southward flowing current beneath
the morthern part of the seasonally reversing Someli
Current; sobteland at moored stations was 7% about 30
is off the Schall count, Its uses amount transport in
the layer 150-600 a manutat to about 3 x 100m/s. The
maderourrent has a pronounced seasonal open of the phase
with the near surface flow, somewhile order in phase
to the someone wind forcing with the spin-up of the
desp-reaching surthern Someli, give after the opact of
the sonthest condent the undercurrent is temporarily
destroyed in the nothbard Schall Basia during June/July,
but is re-agabilished in August. The undercurrent does
the reachil (prest, undercurrent, Indian Open)
(Scophyr, Sec., Greek, Fast, 300433

4711 Chroulation VARIABILITY OF CURRENT, TEMPERATURE, AND BOTTON PERCH ACROSS THE MEST FLORIDA CONTENENTAL SHPLF, WINTER 1941-1942 (1.0. Margarino (A/1) pan Blvd., Tallahasses, Fields, 12301)

undervations are analyzed from four current over smootings deployed on the bread embinated shall the mortings deployed on the bread embinated shall the forthcontern Guif of Heaten from Boxedon 29, 1981, a Pebroary 8, 1982 (71 days), consistent with recet socialing studies, the short of trends too capands with an inertial partial to the alternating up-and-down-cut sympthe scale wind forthing, arong response to 90.3 dyn or 2 along there wind stress (as essayed at the coast) is -20 or a 1 of Cadar Key and -30 or a 1 of coast) is -20 or a 1 of Cadar Key and -30 or a 1 of courth where the shoft infrared (of Capa Smills). Lower layer currents were contarted or and the property of the short of the stress of short of coasts of the pressure field decays of short of folding scale -50 if pressure field decays of short of folding scale -50 if the other of the tide of significant compare sized with the short of the tide singular compare of a vertice variations of the tide singular compare of the short of Observations are qualified from four current with I. Smyllyn, Rost, Green, Paper 1000R

Particles and Fields— Interplanetary Space

3160 Solar Wind Interaction with Moss and Planets
ON YHE NATURE OF THE INTERACTION OF THE SMISS
MAGHETONPHERE WITH THE INTERACTION OF THE SMISS
M. S. Wolff (Space Physics Socion, Jet Propents
Laboratory, Galifornia Institute of TechnologiPasandens, California 91109). B. A Modis
The nature of the interactions between the Smiss
wagnelosphere and Europa, Ganymade, and Galisson
magnelosphere and Europa, Ganymade, and Galisson
examinud. Effects of plasma and storometechnic lerbordeent of each antellite are considered in a disabbordeent of each antellite are considered in a disabthe aurface properties of each of the three senthe aurface properties of each of the three senthe aurface properties of each of the three sentemperatured are used as a basis to culculate smismated) Hyd vapor pressures for each of the three sidlites es a function of matellite latitude and senlites as a function of matellite latitude and sentime. Sublimating H₂O leads to a mat O₂ also splittle such of the three moons, and the effectiveness of mat and the effectiveness of materials and materials and materials are also as a sublimation of the effectiveness of materials and materials are also as a sublimation of the effectiveness of materials and materials are also as a sublimation of the effectiveness of materials and materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of materials are also as a sublimation of the effectiveness of t each of the three moons, and the effections for an almosphere in preventing the Jovian places for online such catellite surface is considered wair a variety of mannetospharic conditions. The adults caning each actalitic surface is consisted warlety of magnetospheric conditions. The affect were also anudical pad used to element the properties of he satellite angents priced were also anudicated pad used to element the properties of he satellite angents placed inside and cutside the jovina placed shall the graphing and conditions appropriate to each stone being definition of local particle, and magnetic fill ger planes, energetic particle, and magnetic fill ger planes. In addition, be obtained to originally over each actalitie surface. In addition, before of charged migrometerorids and public of charged migrometerorids and the satellities. It is perfectly planes and appealment of the satellite surfaces of such considerations and satellites. These efforts suggest the consideration and Sarth and appealment of the satellite surface and compared with evaluable laboratory fight consideration and Sarth and appealment of the satellite surface in the satellites of satellites could get to substantially sufface properties of each of the three mountains and accommonsterorial tombardsend of the satellites and satellites of the satellites (sale) and appealment of the satellites and all satellites and the satellites of the satellites. Jovian magnetosphere).

Particles and Fields— Ionosphere

5530 Nigh Latitude Longspheric Carrests
LONGSPHERIC AND SIEUTLAND CHERRIT DISTRIBUTIONS
LONGSPHERIC FROM THE PROBAT AND MODERATIONS OF AND LABORATORY
LONGSPHERIC FROM THE PROBAT AND MODERATIONS OF AND LABORATORY
MODERATE PROBATION LABORATORY
LONGSPHERIC LABORATORY
LONGSPHERICAL
LONGSPHE

5530 High-latitude ionospheric currents
FRST VMF AURORAL RADAR INTERFLACETIER OBSERVATIONS
J. P. PROVIdekes (School of Electrical Engineering,
Cornell University, Itheca, New York, 14853), W. R.
Swartz, D. T. Partoy, and R. G. Fojer
The radar interferemeter technique first used at the
anguetic equator in Peru is also a very powerful means
for studying auroral plasma instabilities. We present
here the first results, obtained with a 49.92 Milz,
20-25 KM peak power pulsaul radar located in Itheca, NY
42.5° N, 76,4° N, Strong auroral achoes were obtained during several highly active periods. Phase
differences between the signal received on the two
antennas accurately decornine the E-N position, within
the scattering volume, of localized scattering conters,
and changes in this phase determine the corresponding
velocity. The signal Deppier shift describes radial
information on the turbulent structure of the achoing
region and show clearly that different features in the
Deppler power spectrum often represent signals coming
from different locations. If we assume that the radia
and tragaverse phase velocities copresent cal drift
velocities, we can determine full harizontal velocity
vectors from the data and hence the horizontal electric
field, usually with a time resolution of the order of
15-30s,
Goophye, Res. Lett., Paper 3LO330 ophys. Res. Lett., Paper JL0330

for use on surface magnetosster data and recently used by Sausjohann et al. (1979). Data sequired by MAGSAT on two days (n 1979) were analyzed in datail. The first was Dat. 21, 1979, a resonably quiet if K.—a-1 time in the southern healsphere; Mhoreas, the ascend example deals with a rather active time, Nov. 17, 1979 (I K.—15t) in the nurthern heatsphere. Important results of this study include the fact that the Birkelend and alvetrojat current systems are colocated. In addition, were realistic tenospheric current distributions have been developed. The analyses have shown a tendency for triangular rather than constant electrojet current distributions as a function of latitude, which is also consistent with the statistical, uniform Region 1 and Region 2 Birkeland current patterns.

J. Geophys. Res., Blue, Paper 140393

(elds-fougsphere) Theograph Cherent Radar spectra in the auroral f

A THEURY OF COMERRENT RADAR SPECTRA IN THE AUROPAL E REGION

J.-P., bt.-learlice (Center for Atmospheric and Space Sciences, Utah State University, Logan, Utah 84.22), Rejector, Schugel We have developed a theory of auroral radar spectra for conditions under which Farley-Bundson waves should be strongly excited in the high latitude E region. The theory samedae that the unerable waves spend the larger part of their lifettee in a linear growth phase. It also uses a previously emexplored property of the waves, andely that waves with different wave vactor components along the bignetic field have markedly different waves to active and of the special control of the widel have calculated the widels of the special, their mean Boppler shift, and anisotropy. We have studied how the properties of special maisotropy. We have studied how the properties of special presence of ion drifts and neutral winds, by the viectron temperatures and densities, by the meetral stone presence of ion drifts and neutral winds, by the viectron temperatures and densities, by the meetral stone presence of the results awent to be in good qualitative agreement with existing observations.

J. Geophys. Res., Blue, Paper 3A0366

Particles and Fields— Magnetosphere

5733 Magnetic storms
MAGNETIC FIELD LINE RECONNECTION 5, CURRENT DISRUPTIONS
AND DUBLE LAYERS
R.L. Stenzel (Department of Physics, University of
California, Los Angeles, CA, 90024), W. Gatelman
and N. Wild
In a large laboratory plasma-a current sheet is
generated in the process of magnetic field line
reconnection. The stability of the sheet with respect
to local current increases is investigated. When
the current density in the center of the sheet
wroads a critical value, spontaneous local current
disruptions are ubsurved, the current from the center
of the sheet moves out to the sides. Magnetic
flux variations in regions remote from the current
these generate an inductive voltage in the current
thop which drops off inside the plasma in the form of
a potential double layer. This leads to particle
acceleration with velocities much larger than those
expected from the steady state electric fields
in the plasma. The particle beams acquire their
energy on expense of the stored magnetic field energy
of the current system. Beam-plasma instabilities
are generated which dissipate some of the directed
kinetic energy and heat the back ground plasma.
A modal for the machanism of the current disruptions
is formulated. The potential structure leads to jon
expulsion creating a localized density drop. The
associated current drop in an inductive circuit
drives the potential structure, thereby providing
feedback for the disruptive instability. It saturates
at a total current loss upon which the currant
system recovers, and the process repeats randomly,
Similarities and differences to magnetospheric System recovers, and the process repeats randomly. Similarities and differences to magnetospheric substorm phenomena are pointed out. (Reconnection, double layers, current disruptions, particle energization).

J. Geophys. Res., Blue, Paper 180428

5755 Pleama instabilities
OBSERVATIONS OF LNR NOISE
OBSERVATIONS OF LNR NOISE
SOUMHING ROCKET S29 BARILM-GEOS
Hannu E. J. Koakinen (Uppsala lonospharic Observatory,
3-7590 Uppsala, Sweden), Gunnar Holmgren, and Paul M.
Kintner

S-79590 Uppsala, Sweden), Gunnar Holmgren, and Paul M. Kining.

The density fluctuation and alactric field data from the Swedish sounding rocket 529 Barium-GFOS shows a broad electrostatic noise band near the lower hybrid fraquency. The noise is related to the spin of the rocket and extends well below the local lower hybrid resonance frequency. Above the altitude of 300 bm the house shows banded structure roughly organized by the hydrogen cyclotron frequency. Simultaneously with the banded structure a signal near the hydrogen cyclotron frequency is detected. This signal is also spin modulated. The character of the noise is discussed in comparison with the probable free energy sources and it is suggested that the noise is locally generated by the rocket paylohd disturbing the plasma. If this interpretation is correct, we expect plasma wave experiments on other spacecrafts to observe similar phenomena. (Lower hybrid resonance, density fluctuations, free energy source).

J. Geophys. Res., Blue, Paper 3A0126

3799 General (Upetream Energetic Ions)
DEFRINGENCE OF 50 key UPETRIAN TON EVENTS AT IMP 7/8
UPON MAGNETIC FIELD-BOW SHOOK GEOMETRY,
D. G. Mitchell (Applied Physics Leberatory, Johns
Hopkins University, Laurel, Maryland, 20707) and E. G.
Realef septime Oniversity, Leurel, Maryland, 20707) and E. G. Realof

He present the results of a statistical study of 4 years (1972-1976) of 100 7 and 8 observations at ~ 40 R. of 30-200 keV upstramm ion events measured with the MIDI/AP. See the septiment. We find a monotopic increase in the probability of observing spatters particle events with a degreese in the angle (8_m) between the interplanetary magnetic field (100) and the Local shack normal at the point where the 100 interests the bow shock, independent of the length of time of box shock connection (12 in greeso ~ 10 almutes). We find cromply equal probability of observing an event above a given flux from any portion of the bow shock with the same value of 0_{max} a growth time of the 50 ~ 200 keV events of 7 ill almutes, a maximum attainable flow of ~ 2.5 x 105 (cm² e sr) , and a positive norrelation between the probability of exceeding a given thus and the 3-hour K, index. The results leply that the local structure of the how shock in the immediate violatity of the field line connection in the generation process of smagnetic upstress particle events, and that wave-particle interactions produce a salf-throtting mechanies that insist produce a salf-incepting mechanies that limits the samisue flux of long eacaping the upstrame Coreshorts.

1799 General (Upstrame Energatic Lond) 1882-1/IMP-5 ORESEVATIONS OF STRULTAMEOUS UPSTREAM 10F Sverra D. G. Hitchell (Applied Physics Laboratory, Johns Bopkins University, Laurel, Haryland, 20707), E. G. Roelet, T. E. Sandarson, E. Ezinhard and E.-P. Wennel-Propagation of upstress energetic (50-200 keV) forb is analyzed in sixtess events observed simultaneously

ione escaping the upstrama foreshock. . Geophys. Hea., Sive. Paper 340399.

by solid-erate detectors on ISER-) at ~ 200 R and on IMP-B at ~ 35 R from the Earth. Conclusions are based on comparisons of the pitch angle distributions observed at the two spacerait and transformed into the solar wind frame. They are been-like at ISER-), and confined to the autwert besimphers. When IMP-B is furthest from the bow shock they are also usually boam-like, or hamispheric. However, when IMP-B is closer to the bow shock, pancaka-like distributions are observed. This systematic variation in the IMP-B pitch angle distributions delimits a scattering region t 15 R systematic variation in the IMP-B pitch angle distributions delimits a scattering region t 15 R systematic variation in the IMP-B pitch along the interplementary angestic field) which desinates ion propagation, inflances the global distribution of fluxes in the foreshock, and may play a tole in acceleration of the fone, when IMP-B is beyond 1 ~ 15 R₀, the propagation appears to be assentially scatter-free between IMP-B and IMP-B in this is deduced from the absence of sarthward lives at IMP-B as well as the tendency for the spin-averaged fluxes to be comparable at the two spaceraft.

J. Geophys. Fos., Riue, Fapat 1A0198

Physical Properties of

6110 Elosticity, fracture and Flow.

REDROCK FRACTURE PARAMETERS FROM THE INTERPRETATION OF WELL TIDES.

Don R. Bener(Department of Energy, Mines and Resources, Earth Physics Branch, Gravity, Goothermies and Caodynamics Diefeison, I Ubservatory Crescent, Octaws, Ontario, Canada, VIA 073)

This paper presents a theory to account for the main features of the tidal response of an isolated plane fracture as revealed by the vator level in an luter-secting agen berehole. According to the theory, the diurnal and semidiurnal idal constituents (both amplitude and phase) of water level variations depend on the fracture aperture, the orionisting (dlp and strike) and the radial extent of the fracture, and the compressibility of the asperitive. The theory is applied to water level observations made in six open bereholes located in crystalline rock at a rest site on the Chalk River Nuclear Laboratories, Accole Energy of Canada Linted in Chalk River, Ontario, Canada. The orsholes were uncased and intersected sany open fractures so that a definitive test of the theory could not be ande. However, in four cases a fracture model could be found which fitted the observations within the experimental error.

J. Geophys. Res., Red, Paper 1808-29

6110 Electicity, Fracture and Plow STRESS CORPOSION AND CRACY PROPAGATION IN SIGUX QUARTZITE

QUARTITE
Lindamae Peck (Department of Geology and Capphysics,
Yale University, New Haven, Connecticut, 06511)
Water-induced changes in the apount of specific energy
(anergy per unit area) consumed during crack setemation
in Bious quartrite wors determined using the analysis
of linear closkic fracture machanics. Madge-loaded
durble contillator. duble cantilever heatter openmica. Range-losaed double cantilever heat test specimens were cracked to air, water, and equeous solutions of MaON, NCI, NCI and mothanol at rack value(ties of 4.2 x 10° to 4.2 mm/s and temperatures of 17-80°C. Scanning slectron microscopy and the doternination of enhanced permeability were used to detect fracture-induced nicro-machine.

mashlifty were used to detect fracture-induced micro-cracks.

The fracture energy of Sious quartaits in water is 15% lower than that of quartaits tracked in air sambless thus ideas that it is a second or the fracture energy on crack velocity (time) and on the concentration of water in the test environment is characteristic of the occurrence of stress correction as mechanism of enhanced crack growth in the presence of water. Increased crack velocity in quartaite cracked in bot water is evidence of thereally enhanced water-weshoning, but the fracture energy of quartaite cracked in hot air or hot water (40-50°) is 12-13% higher than room temperature values. The increase in fracture energy perature values. The increase in fracture energy upon heating the quartrice is attributed to changes (reduction in residue) stress, existence of thurns attrease) in the not stress field responsible for picrocracking and crack propagation in the quartaite. (Stress of the contraction of the contraction. corrosion, crack propagations.
J. Goophys. Ros., Red. Paper 380412

bild Equations of state

SHOCY COMPRESSION OF DIAMOND CRYSTAL

K. Kondo and T. J. Antenn (Setemotogical Laboratory, Persadena, California 9122)

Two shock wave experiments employing inclined altrors have been carried out to determine the Hugoslot elastic limit (HEL), float shock state at 191 and 217 GPs. and the post-shock state of diamond crystal, which is shock-coapressed along the intermediate direction between the (III) and (110) crystalingraphic axes. The HEL wave has a velocity of 19.9 1 0.3 mm/used and an amplitude of 63 2 28 GPs. An siternate interpretation of the inclined wedge mirror streak record suggests a resp precursor wave and then another HEL value. The markum post-shock state is -1.35 Kg/m², this requit suggests as elastic unloading effect or shock-induced transition to a denser (possibly meralic) phase. (Diamond, phase change, failure), Goophys. Res. Lott., Paper 310262

DISPERION IN TROUBLE FLOW TO PRACTURED GROTHERUAL SYSTEMS

R. N. Horne (Petroleum Engineering Department, Stanford University, California, 94205), F. Rodrigues-G.

Interpretation of tracer tasts is generally beams on the smalpsis of flow through a perous medium. In geothermal reservairs however the principal permanbility arises from fractures, and a perous medium approach is not applicable. The dispersion of tracer material flowing in a fracture is shown to be dominated by molecular diffusion across the fracture - a mechanism known as Taylor dispersion. For typical values of reservoir permanerer, this transverse diffusion will obliterate any transverse tracer concentration gradients caused by the welpoity profile ecross the fracture. The effective dispersivity is Count to be a function of fraccurs sperture, man flow valuacity and moleculer of fraccurs sperture, man flow valuacity and moleculer antificated the two process medium flow. Typically Peclet unbares are antificiated to the between orders I and 100, and field results confirm this range. (Disparsion, tracer, geothermal, fractures)

Geophyn. Res. Lett., Paper 310331 Geophya, Res. Lett., Paper 3L0351

Planetology

631D Atmospheres of planets
DAY AMD FIGHT MODELS OF THE VENUS THEMOSPHERE
SIT Heasts (Lumar and Planetsry Laboratory, University
Of Arlaons, Tutson, Arizona 63721), B.W. Sunten, B.R.

of Arisons, Totson, Arisons 657213, R.R. Susten, U.S. Sowell

A model strosphere of Vanus for slittudes between 100 and 178 he is presented for the day soe aight sidesDensities of 602, 60, 0, 82 He, and 02 on the day side, for 8 and 16 hours lokal time, are obtained by simultaneous soultions of confinity equations. These equations couple ionospheric and neutral chealitry, and the transport processes of molecular bad day diffusion. Photodissociation and photoionization 2 coefficients are presented to facilitate the ionosporation of chemistry into circulation models of the Vanus almosphere. Midnight deposition of Co., 60, 6, 8, 8, and 8 are derived from integration of the continuity equations, subject to specified fluxes. The continuity equations, subject to specified fluxes with the observed sirglow of HD and Os('a). The homopause of Vanus is located near 131 be by both the day and night sides.

J. Geophys. Res., Blue, Paper 3A0196

I. Geophys. Res., Blue, Paper JA0196
6510 Atmospheres of Finance
10'S ATMOSPHERE: PRESSUE CONTROL ST REGOLITH COLD
TRAPPIED AND SUPPACE VEHICUS
Dannis L. Marson and Dongias B. Resh (Forth and Space
Sciences Division, Jet Propulsion Laboratory, California Institute of Technology, Tanadama, CA, 9,109)
In a second for the beast pressure of Lo's stmosphere is developed. This model takes into seconds the
praviously ignored fact that much of lo's surface had
praviously ignored fact that much of lo's surface had
praviously ignored fact that much of lo's surface had
praviously ignored fact that much of lo's surface had
praviously ignored fact that much of lo's surface had
praviously in the second face of the portion of the present of second property
allows afficient, subsurface had trapping of stmospheric goodel, are
sured by person to local durface temperature vis. the
SO, vapor-pressure equilibries curve. Sear lo's solve
soler point the pressure is equilibries with a surface
BO, frost deposit is - [O'] bar. Perous murface
models of the type of develop invoke equilibries with

the colder, subsurface permafaget (at ~ 3 cm depth) and yield personnes of ~ 10⁻¹ bar. The subsurface cold trapping endel explaint same but now all observations relevant to lo's atmosphere. The new basel pressure model and the earlier surface frost equilibrium model, when taken together, provide, respectively, lower and upper limits on the basel 50, pressure of lo's atmosphere. If o strongburn, lo surface, suffurdicted, respectively, lower and upper limits on the basel 50, pressure of lower than the lower tha

Seismology

6940 Phanomeus (Serthquate prediction)
RADON AND TILT ANYMALIES DETECTED REPORE THE IMPISIA
(SOUTH ITALT) KARTROGARE OF MOVEMBER 23, 1980 AT GREAT
DISTANCES FROM THE EPICENTER.
L. Allegel, F. Bella. C. Bella Monica, A. Ermini,
S. Improta, V. Sgrigna and F.F. Bingi (letitoto di Fimica, Universita' di Roma, P.1a A. Moro 2, 00185 Rome,
Italy)

Italy)
A strong earthquake (N-5.5) occurred on November 23, 1986 is South Italy (Irpinia region) at hypocantral depth ranging between 10 Km and 18 Km. Anomalous variations in groundwater radum contout, 23% and 170% compared with the background level, were detected at Ross and Biail sites before the earthquake. A tilt snownly, 20 red in amplitude, was observed at L'Aquila site. All the stations were located at distances of about 250 Km from the epicanter. The snownlows duration, for both the examined parameters, lasted five-six months. We suggest to interpret such snownline as pracursors of the (Tpinia serthquake. (Radon, tilt, earthquake prediction).

Geophys. Res. Lett., Paper 310307

6950 Sciency spectrum, space, and time distriquency spectrum, space, and time distri-bution)
BREAKING OF A SINGLE ASPERITY: RUPTURE PROCESS AND SETSHIC RADIATION S. Das (Lamont-Obserty Coological Observatory of Columbia University, Pollandes, New York 10964)

The problem of apentaneous shear rupture of a single circular saperity on an infinite fault plane is studied. Initially, the fault plane is heaten everywhere except at a circular saperity. An applied displacement at infinity results in a stress concentration along the bounding edge of the superity. The frictional stress on the broken part of the fault plane is taken to be a constant. Once a point on the especity breaks, the stress there drops to the same value as on the 'main' fault surface. The rupture is started by relaxing the shear stress at a point on the asperity edge and is then allowed to propagate spontaneously using a critical stress lovel fracture criterion. The rupture process is calculated mamprically. It is found that for appartites of constant strength, the rupture first propagates around the edge of the asperity and then inward, a phenomenon best described by the soil-known tere of classic military amenuer— 'the double enviroling pincer supement'. In the appendix, the expressions for the far-field selselo radiation due to the rupture of such an asperity are derived. It is shown that the n-th cartosian component of the far-field displacement at (s,t) for P, SV and Si waves, using the notation of Aki and Richards [1980], is given by

 $u_n(\vec{x},t) = \frac{\mathbf{n}_{nt}}{4\pi\mu\epsilon^2\vec{R}} \int\limits_{S_0}^{T} \tau_{nt}\left(\vec{\xi},t-\frac{\vec{R}-\vec{\xi}\cdot\vec{T}}{\epsilon}\right) dS(\vec{\xi})$

Thus, the far-field pulses can be directly found from the stress-draps on the fault-plane. This formula is also true for 'orack' or 'dislocation' problems. The directivity function D₁₁ for displacement for the aspertty problem is found to be that for the double couple, modified by some factor. In particular, the fault plane is a nodal plane for SV mayor. (Circular asperity, rupture, sales or addition).

J. Geophys. Res., Red, Paper 3R0269

Geomagnetism and Aeronomy

Volume 22. Number 5

Dorman L. I., Katz M. E., Nosov S. F., Fedorov Yu. I., Shakhov B. A. Modulation of monounergetic spectrum of galactic cosmic rays
Percyasiova N. K., Pokrevsky P. E., Slodzhkov Yu. I. Absorption of galactic cosmic rays in the almosphere in the period 1961 to 1980
Samokhin M. V. Estimation of velocitity of a return streaming when the geomagnetic dipole is oriented slong a non-disturbed velocity of solar wind Erukhimov L. M., Kogan L. M., Myasnikov E. N. About a heating mechanism of the origin of ionosphere F-layer irregularities
Kolesnik A. G., Golikov I. A. A three-dimension model of high-latitudinal F region with the account of non-coincidence of geographical and geomagnetic coordinates

coordinates

Maximova N. M., Osipov N. K. A dynamic model of the main types of convection, longitudinal currents and volumetrical structure of the polar ionosphere Izraitel A. G., Tzedilina E. E. Dependence of a equinocallal analytical model of electron concentration of the ionosphere on solar activity.

Koen M. A., Sidorov I. M. The account of non-atationary and non-linear terms in the equations of movement in solving tasks of ionospheric modelling.

Gokov A. M., Martynenko S. I., Misura V. A., Piven L. A., Somov V. G., Federenko Yu. P., Chernogor L. F., Shemet A. S. Profiles of electron concentration in the night highlatitudinal low ionosphere, artificially disturbed by powerful radiowaves

ful radiowaves

Gaillt T. A., Gusev V. D., Ivanov M. I., Perchalina E. O. Investigation of anisotro-teor component of an ionosphere-scattered signal Kazanskaya Yu. B., Lazarev V. I., Tel'tzov M. V., Lipovetzky V. A., Tulinov V. F., Feigin V. M. Characteristics of suroral electrons as observed at the ASE «Me-

teore
Lyatzky V. B., Leontlev S. V. A neutral vortex, induced by an aurora arch
Serebryakov B. E. Investigation of processes in the thermosphere during the mag-

netic disturbances
Kazimirovsky E. S., Schminder R., Kürschner D., Petrukhin V. F. Continental efficies in a wind field in the region of the upper mesopause at middle latitudes
Andreeva L. A., Katasev L. A., Kosenkova A. V., Latov Yu. O., Uvarov V. M.,
Shipilov V. V., Tahuka T. I. A drift of barium clouds and estimation of
some parameters of longitudinal currents as observed on island Heiss
Danilov A. A., Makarov G. A. Dependence of magnetic activity on orientation of
the geomagnetic dipole relatively to IMF

the geomagnetic dipole relatively to IMF

Alperovitch L. S., Drobzhev V. I., Sorokin V. M., Troitzkaya V. A., Fedorovitch G. V. About middle-latitudinal oscillations of the geomagnetic field and their connection with dynamics of processes in the ionosphere. Metelkin E. V., Sorokin V. M., Fedorovich G. V. About the nature of oscillations of a geomagnetic field generated by solar flares. Shirman B. L. Electrojets of explosive phase of a substorm.

Tzirs G. P. About an attempt to expose the statistical effects of the influence of usual radioradiations on variations of polar substorm elements. Barkhalov N. A. Fading of ultralewirequential waves in the transition region Tverskaya L. V., Khoroshava O. V. About the nature of the source of the package of fading long-period geomagnetic pulsations.

Tkachev G. N., Lankis L. K. Some possibilities of the pick out of the influence of geoclectrical irregularities on polarisation of geomagnetic pulsations in middle latitudes.

Zhuraylev V. I., Sidorin A. Ya. About forecast of hindrances in sounding the Earth crust by separate impulses of current.

Earth crust by separate impulses of current

Lovadny V. I., Sagoria A. Ya. About forecast of integrates in sounding the

Earth crust by separate impulses of current

Lovadny V. I., Paylova I. V. Magnetotelluric sounding of horizontally-flaky medium with tensor electroconductivity of layers

Rotanova N. M., Papliashvili N. E., Pushkov A. N. Space-time analysis of 60-year

variations of the field by the data of the world network of observatories

Braginsky S. I. Analytical description of variations of the geomagnetic field for

the 20th century

Brief information

Ponyavin D. 1., Pudovkin M. I. Geoeffectivity of a meridional component of the large-scaled magnetic field of the Sun

Alexanjan T. M., Bednazhevski V. M., Belov A. V., Blokh Ya. I., Dorman L. I.,

Kaminer N. S. Experimental check of theoretical calculation of the geomagnetic cut off rigidity

Sharving K. N. About a possibility of registration of electrons with E. 20 MoB in the region of the Brazii magnetic anomaly.

Vinlizky A. V., Sukhomazova G. I., Tshepkin L. A. About the connection of parameters of F2 and F1 ionospheric layers in the day to day changes.

Serebryakov B. E. About the dependence of parameters of a disturbed middle-intitudinal F2 region of the ionosphere on local time.

Novikov G. G., Tzigankov S. F. About the dependence of the effective diffusion coefficient, defined by radiometeor data on the wave length.

Dvinskikh N. I., Pisarsky V. Yu., Rudneva N. M. About non-stationarity of connection of parameters of solar wind with AE and ap indexes of magnetic activity. activity
Ponomarev V, N. Estimation of the value of density of longitudinal current in Arykov A. A., Lystzky V. B., Mel'tzev Yu. P. The origin of a longitudinal electric field when the magnetospheric plasma is heated
Lystzky V. B., Mel'tzev Yu. P. About a possibility of exitation of geomagnetic pulsations with a radial direction of IMF
Bolshakova O. V., Troitzkaya V. A. Impulse reconnection as a possible source of relegions of its large. of pulsations of ipcl-typo

Kalisher A. L., Kurchashov Yu. P., Trolizkaya V. A., Feigin F. Z. Haavy lone Othin a radiation helt of the Earth as a possible source of geomagnetic pulsations Pc1-2

Krotevitch N. F., Panurovsky V. N., Rutenko A. N. About the measurement of components of the magnetic field of sea disturbance

Kalishir Yu. D., Rozanova T. S. 11-year variation of a west drift of the magnetic sectors of the Farth. Rolanda N. M., Papitashvili N. E., Pushkov A. N. About interprotation of 60-year variations of the geomagnetic field by quasigarmonic process.

Zinger B. Sh. About an analytical solution of the task of electromagnetic induc-

zinger B. 5n. About an analysical solution of the task of electromagnetic induc-tion in an inhomogeneous thin layer.

Barinov G. I. Influence of anizotrophy on direction of residual magnetisation of thin monocrystal layers of ferric oxido.

Mikhailov Yu. M., Fidelis V. V. About the influence of the nearest zone of a irace after a towing magnetometer on a measurement error

127